



**HAL**  
open science

## Energy relaxation time in heavy ion collisions between 27 and 44 mev/u

B. Borderie, M.F. Rivet, P. Box, M. Dakowski, C. Cabot, Y. El Masri, F.  
Hanappe, D. Gardes, D. Jouan, G. Mamane, et al.

► **To cite this version:**

B. Borderie, M.F. Rivet, P. Box, M. Dakowski, C. Cabot, et al.. Energy relaxation time in heavy ion collisions between 27 and 44 mev/u. Zeitschrift für Physik. A, Atoms and Nuclei, 1997, 357, pp.7-8. in2p3-00003594

**HAL Id: in2p3-00003594**

**<https://in2p3.hal.science/in2p3-00003594v1>**

Submitted on 1 Apr 1999

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

12 novembre 1996

CERN LIBRARIES, GENEVA



SCAN-9701126

IPNO-DRE-96-21

***Energy relaxation time in heavy ion collisions  
between 27 and 44 MeV/u***

*B. Borderie, M.F. Rivet, P. Box, M. Dakowski, C. Cabot,  
Y. El Masri, F. Hanappe, D. Gardès, D. Jouan, G. Mamane,  
X. Tarrago, H. Utsunomiya*

To appear in Zeitschrift für Physik A

SW 9704

## Energy relaxation time in heavy ion collisions between 27 and 44 MeV/u

B. Borderie<sup>a</sup>, M. F. Rivet<sup>a</sup>, P. Box<sup>a</sup>, M. Dakowski<sup>a1</sup>, C. Cabot<sup>a</sup>,  
Y. El Masri<sup>b</sup>, F. Hanappe<sup>c</sup>, D. Gardès<sup>a</sup>, D. Jouan<sup>a</sup>, G. Mamane<sup>a2</sup>,  
X. Tarrago<sup>a</sup>, H. Utsunomiya<sup>a3</sup>

<sup>a</sup> *Institut de Physique Nucléaire IN2P3-CNRS, F91406 Orsay Cedex, France*

<sup>b</sup> *Fonds National de la Recherche Scientifique and Institut de Physique Nucléaire,  
UCL, B-1348, Louvain-la-Neuve Belgium*

<sup>c</sup> *Fonds National de la Recherche Scientifique and Université Libre de Bruxelles,  
B-1050 Bruxelles, Belgium*

### Abstract

The excitation energy sharing between the two partners of binary dissipative collisions was derived for the  $^{40}\text{Ar} + ^{\text{nat}}\text{Ag}$  system at 27 and 44 MeV/u. From the eventual equilibration of energy between the two partners, and their separation time obtained from Landau-Vlasov simulations, the energy relaxation time was extracted, and compared with theoretical predictions.

The knowledge of the different time scales involved in nuclear collisions is of crucial importance at energies above the Fermi energy. In this domain the reaction times become very short, possibly of the order of the energy relaxation time; it therefore becomes difficult to clearly distinguish between a collision stage and a deexcitation stage. For binary dissipative collisions, the interaction time extends up to the re-separation of the two partners. When studying the deexcitation of these fragments, it is generally assumed that they have reached thermal equilibrium. A review of the different time scales involved in nuclear collisions can be found in <sup>1</sup>. In binary dissipative collisions one can experimentally derive the excitation energy sharing between the two primary partners. If both have the same excitation energy per nucleon (i.e. the same temperature), then one may assume that the dinuclear system was equilibrated at re-separation; the associated time has to be derived from a dynamical simulation of nuclear collisions. First data using this method were published in <sup>2</sup>. They were obtained from the reactions of  $^{40}\text{Ar}$  on  $^{\text{nat}}\text{Ag}$  at 27 MeV/u. In this note we report on the same system at 44 MeV/u comparing the energy relaxation times thus obtained with theoretical calculations.

The experimental method is based on exclusive measurements, detecting in coincidence the two final partners of the collisions. Experimental details can be found in <sup>3, 4</sup>. The emission angle (with respect to the beam direction) of the light fragment,  $\theta_L$ , was taken as a reference, then at each corresponding value average kinematic quantities related to both fragments were measured (velocities of the

---

<sup>1</sup> *On leave from: Institute of Experimental Physics, 00-681 Warsaw, Poland*

<sup>2</sup> *Present address: CUEPE, Université de Genève, 1231 Conches, Switzerland*

<sup>3</sup> *Present address: Konan University, Higashinada, Kobe 658 Japan*

light and heavy fragments,  $V_L, V_H$ , angle of the heavy fragment,  $\theta_H$ ). The method used to derive these quantities is described in <sup>5</sup>, and the results in <sup>6</sup>. These data suggest the presence of an abundant preequilibrium emission, which it was difficult to separate, experimentally, from emission by the light fragment. Therefore the contribution of preequilibrium particles (in mass  $A_{PE}$  and energy  $E_{PE}$ ) was taken from Landau-Vlasov dynamical simulations, which were shown in <sup>6</sup> to correctly describe the dynamics of the peripheral and semi-central collisions. Then the primary masses of the light partners of the collision are calculated from the dynamics following:

$$A'_L = \frac{A_{TOT} - A_{PE}}{\sin \theta_L} \times \left( \frac{\langle V_L \rangle}{\sin \langle \theta_H \rangle} + \frac{\langle V_H \rangle}{\sin \theta_L} \right)^{-1} \langle V_H \rangle \quad (1)$$

The subscripts L (H) refer to the light (heavy) partner of the collision, and the prime(') stands for the primary light fragment;  $A_{TOT}$  is the sum of the projectile and target masses. Equation (1) just expresses the conservation of perpendicular linear momentum. The primary mass of the heavy partner follows from total mass conservation. The ratio of the excitation energy of the light fragment to the total excitation energy is taken equal to the ratio of the mass loss of this fragment (primary - measured) to the total mass loss. The calculation was performed at two different angles and for different final Z values of the light fragment. In Fig 1 the excitation energy ratios are plotted as a function of the detected atomic number of the light fragment; for completeness, the data previously obtained at 27 MeV/u are also displayed in Fig 1. The errors bars come mainly from uncertainties in determining the most probable velocity ( $\langle V_H \rangle$ ) and angle ( $\langle \theta_H \rangle$ ) of the heavy partner.

Solid lines in the figure indicate the expected behaviour assuming equal temperatures. It can be concluded from this figure that the equilibrium between the two partners of the collision is far from being reached at 15° at 27 MeV/u but is achieved in the three others cases. This clearly fixes upper limits for the energy equilibration time, namely the re-separation time between the two fragments, which can be derived from the dynamical simulations <sup>6</sup>. This needs a link between the experiment and the calculation, which can be done through the emission angle of the light fragment recalling the strong correlation between this angle and the impact parameter. Results are given in table 1, which restrict the energy relaxation time to the range 100-180 fm/c ( $\leq 125$  fm/c) for an incident energy of 27 (44) MeV/u .

Theoretical calculations of the energy relaxation time have been performed for many years. Calculations in infinite nuclear matter predict that this time decreases by a factor of 5 when the incident energy increases from 20 to 100 MeV/u <sup>7, 8, 9</sup>. In these papers the relaxation is assumed to be essentially due to nucleon-nucleon collisions, and the large variations observed from one calculation to another <sup>1</sup> are due to the different values adopted for the nucleon-nucleon cross section. For finite nuclei, the relaxation time has to be derived from dynamical simulations, and it




Figure 1: Ratio of the excitation energy of the light partner to the total excitation energy for different atomic numbers of the detected light fragment. Data were obtained at two incident energies, and at two detection angles of the light fragment at each energy

$E_{inc}$ (MeV/u)	$\theta_{LF}$	$b$ (fm)	$t_{sep}$ (fm/c)
27	$15^\circ$	8	100
27	$30^\circ$	6	180
44	$10^\circ$	5	125
44	$20^\circ$	3	125

Table 1:

is generally related to the variation of the total quadrupole momentum, which decreases exponentially with time ( $Q_2(t) = Q_2(t=0)\exp(-t/\tau_Q)$ ). For a light system such as Ca+Ca, and the model of Cassing<sup>9</sup>, the value of  $\tau$  is  $\sim 10$  fm/c and is found to be independent of the bombarding energy. For the  $^{40}\text{Ar} + ^{nat}\text{Ag}$  system and the Landau-Vlasov simulations of<sup>6</sup>, a higher value of  $\tau_Q$  is obtained, slightly decreasing when the incident energy increases, from 28 fm/c at 27 MeV/u to 18-20 fm/c at 44 and 57 MeV/u. Moreover in this framework the relaxation time was also found to be independent of the total mass of the system, as similar  $\tau_Q$  values were found

for the Ar+Al <sup>10</sup> and Gd+U <sup>11</sup> systems, both studied at 35 MeV/u . In ref <sup>12</sup>, the authors of the Landau-Vlasov simulation define a local equilibrium time, related to the relaxation of the Fermi bisphere, which is 25% lower than  $\tau_Q$ , but has the same behaviour as a function of the incident energy.

Full equilibration of energy should be reached around  $t \sim 3-4$  times  $\tau_Q$ , i.e. 100 fm/c for Ar+Ag reactions. In this sense, and relying consistently on the Landau-Vlasov simulations to derive both the re-separation time and the equilibration time, the "experimental" energy relaxation times derived from table 1 are in good agreement (within 25%) with the calculated values.

## References

1. B. Borderie, Ann. Phys. Fr. 17 (1992) 349
2. B. Borderie et al, Z. Phys 338 (1991) 369
3. M.F. Rivet et al, Proc. of the XXXI Int. Winter Meeting on Nuclear Physics, Bormio, (1993) - (Ricerca Scientifica ed Educazione Permanente, Milano, 1993) p92
4. P. Box et al, Proc. Second European Biennial Conference on Nuclear Physics, Megève, (1993) - (World Scientific, ed by D. Guinet, 1995) p 237
5. D. Jouan et al., Z. Phys. A-Hadrons and Nuclei 340 (1991) 63
6. F. Haddad et al, Z. Phys A354 (1996) 321
7. G.F. Bertsch, Z. Phys. A289 (1978) 103
8. C. Toepffer and Cheuk-Yin Wong, Phys. Rev. C25 (1982) 1019
9. W. Cassing, Z. Phys. A327 (1987) 447
10. C. Grégoire et al, Nucl. Phys. A465 (1987) 317
11. B. Borderie et al, Phys. Lett. B302 (1993) 15
12. P. Abgrall et al, Phys. Rev. C49 (1994) 1040