



HAL
open science

Dispersion relation results for VCS at JLab

G. Laveissiere

► **To cite this version:**

G. Laveissiere. Dispersion relation results for VCS at JLab. Compton Scattering from Low to High Momentum Transfer, Mar 2003, Trento, Italy. pp.1-16. in2p3-00012877

HAL Id: in2p3-00012877

<https://in2p3.hal.science/in2p3-00012877v1>

Submitted on 19 Jun 2003

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.


L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Virtual Compton Scattering @ Jefferson Lab

Extraction of Proton Generalized Polarisabilities
at $Q^2=1.0$ and 1.9 GeV^2
using a Dispersion Relations formalism

Compton Scattering from low to high momentum transfer
Trento, Italy, April 2003

Electroproduction of photons experiment: analysis below and above π^0 production threshold



$$d\sigma_{exp}^0 = \frac{N_{exp}}{N_{sim}} \times \frac{L_{sim}}{L_{exp}} \times d\sigma_{sim}^0.$$

We need an accurate Monte Carlo simulation in order to obtain the experimental cross section.


First order Generalized Polarisabilities are not valid in the Δ region.

Low E_γ	Around π^0 threshold	Δ region
Bethe–Heitler + Born	1 st Order GP	Dispersion Relations


We need a new model to describe the cross sections in the resonance region.

Proton Generalized Polarisabilities extraction above π^0 production threshold

Analyticity
+
Crossing Rules
+
Unitarity




**Dispersion
Relations**



Non-Born
Amplitudes
 \Leftrightarrow
Generalized
Polarisabilities

Intermediate States πN Amplitudes



This formalism has 2 free parameters: the 2 generalized polarisabilities

$$\alpha(Q^2) \text{ and } \beta(Q^2)$$

\Rightarrow valid until the 2 pions production threshold

\Rightarrow related to the structure functions $P_{LL} - P_{TT}/\epsilon$ and P_{LT}

B. Pasquini, M. Gorchtein, D. Drechsel, A. Metz and M. Vanderhaeghen,
Eur. Phys. J. A11 (2001) 185


Jlab E93050

Absolute cross sections measurement


Coincidence e/p

Need good resolution for γ/π^0 separation


Normalization controlled at low E_γ




Missing mass (MeV^2)



Vertex Reconstruction (mm)



Coincidence Time (ns)



Analysis

BELOW π^0 THRESHOLD

Low Energy Theorem Analysis \longrightarrow VCS Structure functions

$$Q^2 = 0.923 \text{ GeV}^2 \quad \varepsilon = 0.950$$

$$Q^2 = 1.760 \text{ GeV}^2 \quad \varepsilon = 0.879$$

$$\mathbf{P}_{LL} - \mathbf{P}_{TT}/\varepsilon, \mathbf{P}_{LT}$$

ABOVE π^0 THRESHOLD

Dispersion Relations Analysis \longrightarrow 2 DR Parameters

$$Q^2 = 0.923 \text{ GeV}^2 \quad \varepsilon = 0.950$$

$$Q^2 = 1.760 \text{ GeV}^2 \quad \varepsilon = 0.879$$

$\Lambda_\alpha, \Lambda_\beta$ related to α and β via a dipole form:

$$\alpha(Q^2) - \alpha^{\pi N}(Q^2) = \frac{\alpha - \alpha^{\pi N}}{(1 + Q^2 / \Lambda_\alpha^2)^2}$$

$\Delta(1232)$ REGION

Dispersion Relations Analysis \longrightarrow 2 DR Parameters



$$Q^2 = 0.923 \text{ GeV}^2 \quad \varepsilon = 0.950$$

$$\Lambda_\alpha, \Lambda_\beta$$




Cross sections at $Q^2 = 0.923 \text{ GeV}^2$ above π^0 production threshold

Lepton Plane

40° Out-Of-Plane





Dispersion relations fit at $Q^2 = 0.923 \text{ GeV}^2$ above π^0 production threshold






Cross Sections at $Q^2 = 1.760 \text{ GeV}^2$ above π^0 production threshold

Lepton Plane


40° Out-Of-Plane






Dispersion relations fit at $Q^2 = 1.760 \text{ GeV}^2$ above π^0 production threshold



Cross sections at $Q^2 = 0.923 \text{ GeV}^2$ in the Δ resonance region




Dispersion relations fit at $Q^2 = 0.923 \text{ GeV}^2$ in the $\Delta(1232)$ region



Comparison of structure functions

3 different analysis


$Q^2 = 0.923$
 GeV^2



Comparison of structure functions

LEX and DR analysis


$Q^2 = 1.760$
 GeV^2





Comparison of dispersion relations parameters

3 different analysis

Dipole form is
pretty good !!



Summary



proton EM Form factors = parametrization of Brash et al.


exp. point at $Q^2=0.00$ / (RCS) V.Olmos de Leon et al., Eur.Phys.J.A10 (2001) 207

exp. point at $Q^2=0.33$ / VCS (Mami) J.Roche et al, Phys.Rev.Lett.85 (2000) 798

exp. point at $Q^2=0.92$ and 1.76 / VCS JLab E93050 LEX analysis

exp. point at $Q^2=0.92, 0.92$ and 1.76 / VCS JLab E93050 DR analysis

VCS / RCS Comparison



Cross sections of the same order (Q^2), Identical Resonances VCS follows « scaling »?