



HAL
open science

Measurement of triple gauge-boson couplings in ALEPH

R. Bruneliere

► **To cite this version:**

R. Bruneliere. Measurement of triple gauge-boson couplings in ALEPH. Lake Louise Winter Institute, Feb 2002, Alberta, Canada. in2p3-00014174

HAL Id: in2p3-00014174

<https://in2p3.hal.science/in2p3-00014174v1>

Submitted on 25 Nov 2003

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

MEASUREMENT OF TRIPLE GAUGE-BOSON COUPLINGS IN ALEPH

R. BRUNELIÈRE

LAPP Chemin de Bellevue 74940 Annecy-le-Vieux, FRANCE

E-mail: brunelie@lapp.in2p3.fr

The triple gauge-boson couplings involving the W are determined using data samples collected with the ALEPH detector at mean centre-of-mass energies between 183 GeV and 208 GeV, corresponding to an integrated luminosity of 683.6 pb^{-1} . The triple gauge-boson couplings, Δg_1^Z , $\Delta \kappa_\gamma$ and λ_γ are measured using an optimal observable analysis of W -pair event topology. Results from single- W and single- γ production are included. The three couplings are measured individually assuming the two other couplings to be fixed at zero (Standard Model value). In addition, we use W -pair events to set limits on the C- and P-violating couplings g_4^V , g_5^V , $\tilde{\kappa}_V$, and $\tilde{\lambda}_V$, where V denotes either γ or Z . No deviations from the Standard Model expectations are observed.

1 Introduction

Triple Gauge-Boson Couplings can be separated in two main categories: vertices which involves only neutral bosons Z or γ and vertices with charged bosons W . The former $VZ\gamma$ and VZZ ($V = Z^*/\gamma^*$) are not present in the standard model at tree level. They are searched at LEP2 in the production of ZZ pairs and in $Z\gamma$ events¹ and won't be treated in the following note. The latter VW^+W^- can be found in three different physics processes at LEP2: WW pair production, single W production and single γ production. The VW^+W^- vertices are a direct consequence of the $SU(2)_L \times U(1)_Y$ gauge theory and are present in the standard model at Born level. Their study represents a fundamental test of the non-Abelian nature of the Standard Model. The most general Lorentz invariant parametrisation of the γW^+W^- and ZW^+W^- vertices can be described by 14 independent complex couplings^{2,3}, 7 for each vertex: g_1^V , g_4^V , g_5^V , κ_V , λ_V , $\tilde{\kappa}_V$ and $\tilde{\lambda}_V$, where V denotes either γ or Z . Assuming electromagnetic gauge invariance, C- and P-conservation, the set of 14 couplings can be reduced to 5 parameters: g_1^Z , κ_γ , κ_Z , λ_γ and λ_Z , with Standard Model values $g_1^Z = \kappa_Z = \kappa_\gamma = 1$ and $\lambda_Z = \lambda_\gamma = 0$. Finally, local $SU(2)_L \times U(1)_Y$ gauge invariance introduces the constraints:

$$\Delta \kappa_Z = -\Delta \kappa_\gamma \tan^2 \theta_W + \Delta g_1^Z, \quad (1)$$

$$\lambda_Z = \lambda_\gamma, \quad (2)$$

where Δ denotes the deviation of the respective quantity from its non-zero Standard Model value, and θ_W is the weak mixing angle. Hence, only three parameters remain: Δg_1^Z , $\Delta \kappa_\gamma$, and λ_γ ³. This paper presents preliminary results on the three couplings Δg_1^Z , $\Delta \kappa_\gamma$, and λ_γ using all ALEPH data, but also updates the results from single-parameter fits to the unconstrained real and imaginary parts of the 8 C- or P-violating couplings ⁴.

2 The ALEPH Detector

A detailed description of the ALEPH detector may be found in ^{5,6}. The central part of the ALEPH detector is dedicated to the reconstruction of the trajectories of charged particles. Following a charged particle from the interaction point outwards, the trajectory is measured by a two-layer silicon strip vertex detector (VDET), a cylindrical drift chamber (ITC) and a large time projection chamber (TPC). They are immersed in a 1.5 T axial field and combined, they measure charged tracks with a momentum resolution of $\delta p_T/p_T = 6 \times 10^{-4} p_T \oplus 0.005$. Photons and electrons are identified in the electromagnetic calorimeter (ECAL), situated between the TPC and the coil. It yields a relative energy resolution of $0.18/\sqrt{E} \oplus 0.009$. The iron return yoke is equipped of 23 layers of streamer tubes and forms the hadron calorimeter (HCAL). Combined with ECAL it provides a relative energy resolution of charged and neutral hadrons of $0.85/\sqrt{E}$. Muons are distinguished from hadrons by their distinct pattern in HCAL and by the muon chambers outside the HCAL.

3 Event selection and kinematic reconstruction

The selection of W^+W^- candidates is highly dependent of the event topology. Selected events are exclusively classified with the following order of priority: $evq\bar{q}$, $\mu\nu q\bar{q}$, $\tau\nu q\bar{q}$, $q\bar{q}q\bar{q}$, and $l\nu l\nu$ according to the way each W is decaying ⁷.

3.1 $W^+W^- \rightarrow evq\bar{q}$ and $W^+W^- \rightarrow \mu\nu q\bar{q}$ events

Semileptonic events are selected such that they contain a high energy lepton candidate and two jets. The selection of the lepton is using track isolation. The DURHAM-PE clustering algorithm is then applied to all objects that are not used to construct the lepton four-momentum, and these are forced in two jets. After this preselection, the probability for the event being signal is determined using a Neural Networks based on the momentum of the lepton, the total missing transverse momentum and the lepton isolation. To improve

the resolution on the reconstructed four-momenta of the W decay products, the events are subjected to a kinematic fit. Typical efficiency of semileptonic channels is roughly 80% for a purity of 95%.

3.2 $W^+W^- \rightarrow \tau\nu q\bar{q}$ events

Selection of semileptonic events with a τ is slightly different from the previous case, since τ lepton can decay into hadrons. In that case, an iterative method is applied to search for the most isolated jet amongst the jets resulting from a low y_{cut} value of $0.75 \text{ GeV}/c^2$ with the JADE algorithm. To improve the resolution of the angular observables a kinematic fit is performed where the direction of the τ is approximated by its visible decay products. Efficiency of $\tau\nu q\bar{q}$ selection is about 54% and purity is roughly 76%.

3.3 $W^+W^- \rightarrow q\bar{q}q\bar{q}$ events

Fully hadronic events are identified by having four separated jets contained in the detector. To extract the hadronic W^+W^- signal with high purity ($\simeq 85\%$) and efficiency ($\simeq 80\%$), the selection is based on the output of a neural network based on 14 variables ⁷. Moreover, for the hadronic W^+W^- events the reconstruction of the relevant information is more complicated since there is no clean signature of the W^- direction nor any information of the particle flavors in either W systems. In this case the four jets can be paired in three different ways. To select the best pairing with a purity of 75%, a 6-constraint kinematic fit is applied to all three possible pairings. To assign a jet pair to the W^+ or W^- , a jet charge algorithm is used. The jet charge is obtained from the pseudorapidity-weighted charge for the particles forming a jet.

3.4 $W^+W^- \rightarrow l\nu l\nu$ events

The efficient of the selection of purely leptonic events is low 27%, but with a high purity 97%. Selection is based on missing transverse momentum, missing mass and kinematic properties of the lepton candidates. For purely leptonic W^+W^- events the momenta of the two neutrinos are unknown. However in the absence of Initial State Radiation and neglecting the W width, the constraint that the two $l\nu$ systems should have the W mass in combination with the usual four-momentum conservation allows a reconstruction of the neutrino momenta.

4 Determination of the TGCs

4.1 Coupling extraction

All the information on the couplings is contained in a five angles distribution and the electric charge of the fermions³. Since triple gauge-boson couplings contribute only linearly to the amplitude of the considered processes, the differential cross section can be expanded in these couplings g_i as a second order polynomial

$$\frac{d\sigma}{d\Omega} = S_0(\Omega) + \sum_i S_{1,i}(\Omega)g_i + \sum_{ij} S_{2,ij}(\Omega)g_i g_j. \quad (3)$$

Couplings estimators are then extracted with an optimal observable method⁸. The general idea of optimal observables is to extract the couplings g_i by measuring the mean values \overline{OO} of distributions of suitably defined observables OO . In addition to the total number of selected events, the two optimal observables builded are

$$OO_{1,i}(\Omega) = \frac{S_{1,i}(\Omega)}{S_0(\Omega)} \text{ and } OO_{2,ij}(\Omega) = \frac{S_{2,ij}(\Omega)}{S_0(\Omega)}. \quad (4)$$

With only Gaussian distributed variables, likelihood function is then equivalent to a least square method. In particular, including systematic errors is easy.

4.2 Systematic uncertainties

A complete list of all the systematics errors computed can be found in⁴. The main sources are coming from uncertainties on the Monte Carlo model used to generate the W^+W^- signal. They were estimated as the effect of higher order correction for these results, but a reduction of the error size is foreseen for the final results⁹. The other main systematic uncertainties are due to fragmentation or final state interactions not perfectly modeled, i.e. Color Reconnection and Bose-Einstein effects.

5 Results and conclusion

The triple gauge-boson couplings have been measured using W-pair events at all LEP2 energies. Combining with the ALEPH measurement from single-W production and single- γ production, the three couplings Δg_1^Z , $\Delta \kappa_\gamma$ and λ_γ have been measured individually. The results are

$$\Delta g_1^Z = 0.015_{-0.032}^{+0.035}$$

$$\Delta\kappa_\gamma = -0.020_{-0.072}^{+0.078}$$

$$\lambda_\gamma = -0.001_{-0.031}^{+0.034},$$

where the error includes systematic uncertainties. The corresponding 95% confidence level limits,

$$-0.048 < \Delta g_1^Z < 0.080$$

$$-0.164 < \Delta\kappa_\gamma < 0.132$$

$$-0.059 < \lambda_\gamma < 0.065,$$

are in good agreement with the Standard Model expectation.

In addition, W-pair events were used to set limits on the C- or P-violating couplings g_4^V , g_5^V , $\tilde{\kappa}_V$ and $\tilde{\lambda}_V$, where V denotes either γ or Z . Results are shown on Table 1.

Table 1. Combined results for the real and imaginary parts of the C- or P- violating couplings from W-pair production at 183-208 GeV. Each coupling is determined setting all other couplings to their Standard Model value. The error includes systematic uncertainties. The corresponding 95% confidence intervals are listed in the last column.

Coupling	Fit result	95% confidence limits
$Re(\tilde{\kappa}_\gamma)$	$-0.027_{-0.111}^{+0.114}$	[-0.236, 0.191]
$Re(\tilde{\lambda}_\gamma)$	$0.001_{-0.090}^{+0.090}$	[-0.170, 0.173]
$Re(\tilde{\kappa}_Z)$	$-0.006_{-0.060}^{+0.060}$	[-0.123, 0.111]
$Re(\tilde{\lambda}_Z)$	$-0.004_{-0.048}^{+0.048}$	[-0.096, 0.089]
$Re(g_4^\gamma)$	$0.116_{-0.158}^{+0.156}$	[-0.193, 0.418]
$Re(g_5^\gamma)$	$-0.186_{-0.189}^{+0.188}$	[-0.557, 0.181]
$Re(g_4^Z)$	$0.103_{-0.120}^{+0.119}$	[-0.134, 0.334]
$Re(g_5^Z)$	$-0.130_{-0.138}^{+0.138}$	[-0.400, 0.141]
$Im(\tilde{\kappa}_\gamma)$	$-0.022_{-0.057}^{+0.057}$	[-0.134, 0.090]
$Im(\tilde{\lambda}_\gamma)$	$0.037_{-0.046}^{+0.045}$	[-0.053, 0.125]
$Im(\tilde{\kappa}_Z)$	$-0.045_{-0.037}^{+0.037}$	[-0.116, 0.027]
$Im(\tilde{\lambda}_Z)$	$0.043_{-0.029}^{+0.029}$	[-0.015, 0.100]
$Im(g_4^\gamma)$	$0.286_{-0.132}^{+0.130}$	[0.027, 0.539]
$Im(g_5^\gamma)$	$-0.180_{-0.216}^{+0.219}$	[-0.600, 0.251]
$Im(g_4^Z)$	$0.167_{-0.082}^{+0.082}$	[0.005, 0.326]
$Im(g_5^Z)$	$-0.089_{-0.142}^{+0.142}$	[-0.366, 0.190]

References

1. ALEPH Collaboration, *Limits on anomalous neutral gauge couplings using data from ZZ and Z γ production between 183-208 GeV*, contributed paper for LP01 and EPS HEP 2001, ALEPH 2001/061, CONF 2000/041.
2. K. Hagiwara, R. D. Peccei, D. Zeppenfeld and K. Hikasa, *Nucl. Phys. B* **282**, 253 (1987).
3. G. Gounaris, J.-L. Kneur and D. Zeppenfeld, from *Physics at LEP2*, CERN 96-01 p. 525, editors G. Altarelli, T. Sjöstrand and F. Zwirner.
4. ALEPH Collaboration, *Measurement of Triple Gauge-Boson Couplings in e^+e^- collisions up to 208 GeV*, contributed paper for LP01 and EPS HEP 2001, ALEPH 2001/060, CONF 2000/040.
5. ALEPH Collaboration, *ALEPH: A detector for Electron-Positron Annihilations at LEP*, *Nucl. Instrum. Methods A* **294**, 121 (1990).
6. ALEPH Collaboration, *Performance of the ALEPH detector at LEP*, *Nucl. Instrum. Methods A* **360**, 481 (1995).
7. ALEPH Collaboration, *Measurement of W-pair production and W branching ratios in e^+e^- collisions up to 208 GeV*, contributed paper to the 2001 Winter Conferences, ALEPH 2001-013, CONF 2001-010.
8. M. Diehl and O. Nachtmann, *Z. Phys. C* **62**, 397 (1994).
9. R. Brunelière *et al*, hep-ph/0201304, CERN-TH-2001-274.