



HAL
open science

alpha decay half-lives of new superheavy nuclei within a generalized liquid drop model

Hongfei Zhang, Wei Zuo, Junqing Li, Guy Royer

► **To cite this version:**

Hongfei Zhang, Wei Zuo, Junqing Li, Guy Royer. alpha decay half-lives of new superheavy nuclei within a generalized liquid drop model. *Physical Review C*, 2006, 74, pp.017304. 10.1103/PhysRevC.74.017304 . in2p3-00087636

HAL Id: in2p3-00087636

<https://in2p3.hal.science/in2p3-00087636v1>

Submitted on 26 Jul 2006

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

α decay half-lives of new superheavy nuclei within a generalized liquid drop model

Hongfei Zhang^{1,2,3,*}, Wei Zuo^{1,2}, Junqing Li^{1,2}, and G. Royer⁴

¹*Institute of Modern Physics, Chinese Academy of Science, Lanzhou 730000, China*

²*Research Center of Nuclear Theory of Laboratory of Heavy Ion Accelerator of Lanzhou, Lanzhou 730000, China*

³*Graduate school of Chinese Academy of Sciences, Beijing 100039, China and*

⁴*Laboratoire Subatech, UMR: IN2P3/CNRS-Université-Ecole des Mines 4 rue A. Kastler, 44307 Nantes Cedex 03, France*

(Dated: July 27, 2006)

The α decay half-lives of the recently produced isotopes of the 112, 114, 116 and 118 nuclei and decay products have been calculated in the quasi-molecular shape path using the experimental Q_α value and a Generalized Liquid Drop Model including the proximity effects between nucleons in the neck or the gap between the nascent fragments. Reasonable estimates are obtained for the observed α decay half-lives. The results are compared with calculations using the Density-Dependent M3Y effective interaction and the Viola-Seaborg-Sobiczewski formulae. Generalized Liquid Drop Model predictions are provided for the α decay half-lives of other superheavy nuclei using the Finite Range Droplet Model Q_α and compared with the values derived from the VSS formulae.

PACS numbers: 27.90.+b, 23.60.+e, 21.10.Tg

The synthesis of superheavy elements has advanced using both cold [1] and warm fusion reactions. Recently [2, 3, 4], isotopes of the elements 112, 114, 116 and 118 have been produced in fusion-evaporation reactions at low excitation energies by irradiations of the ^{233,238}U, ²⁴²Pu, ²⁴⁸Cm and ²⁴⁹Cf targets with ⁴⁸Ca beams. The main decay mode is the α emission and the α decay energies and half-lives of fourteen new α decaying nuclei have been measured. Some questions have been raised [5] about these superheavy element findings. In similar sophisticated experiments at other places [6, 7] the α cascades were not observed.

The pure Coulomb barrier sharply peaked at the touching point does not allow to determine correctly the partial α decay half-lives. It is probable that the α decay takes place in the quasi-molecular shape path where the nucleon-nucleon forces act strongly during the formation of the neck between the nascent fragments and after the separation and a proximity energy term must be added in the usual development of the liquid-drop model [8]. The generalized liquid drop model (GLDM) which includes such a proximity energy term has allowed to describe the fusion [9], fission [10], light nucleus [11] and α emission [12, 13] processes.

The purpose of this work is to determine the partial α decay half-lives of these superheavy elements within this GLDM from the experimental Q_α values using the WKB approximation and to compare with the experimental data and the calculations with the Density-Dependent M3Y (DDM3Y) effective interaction [14] and the Viola-Seaborg formulae with Sobiczewski constants (VSS) [15]. Finally predictions within the GLDM and VSS formulae are given for the partial α decay half-lives of the still non observed superheavy nuclei ranging from Sg to $Z = 120$.

For a deformed nucleus, the macroscopic GLDM en-

ergy is defined as [9].

$$E = E_V + E_S + E_C + E_{\text{Rot}} + E_{\text{Prox}}. \quad (1)$$

When the nuclei are separated:

$$E_V = -15.494 [(1 - 1.8I_1^2)A_1 + (1 - 1.8I_2^2)A_2] \text{ MeV}, \quad (2)$$

$$E_S = 17.9439 [(1 - 2.6I_1^2)A_1^{2/3} + (1 - 2.6I_2^2)A_2^{2/3}] \text{ MeV}, \quad (3)$$

$$E_C = 0.6e^2 Z_1^2/R_1 + 0.6e^2 Z_2^2/R_2 + e^2 Z_1 Z_2/r, \quad (4)$$

where A_i , Z_i , R_i and I_i are the mass number, charge number, radii and relative neutron excesses of the two nuclei. r is the distance between the mass centres. The radii R_i are given by [8]:

$$R_i = (1.28A_i^{1/3} - 0.76 + 0.8A_i^{-1/3}) \text{ fm}. \quad (5)$$

This formula allows to follow the experimentally observed increase of the ratio $r_i = R_i/A_i^{1/3}$ with the mass; for example, $r_0 = 1.13$ fm for ⁴⁸Ca and $r_0 = 1.18$ fm for ²⁴⁸Cm.

For one-body shapes, the surface and Coulomb energies are defined as:

$$E_S = 17.9439(1 - 2.6I^2)A^{2/3}(S/4\pi R_0^2) \text{ MeV}, \quad (6)$$

$$E_C = 0.6e^2(Z^2/R_0) \times 0.5 \int (V(\theta)/V_0)(R(\theta)/R_0)^3 \sin \theta d\theta. \quad (7)$$

S is the surface of the one-body deformed nucleus. $V(\theta)$ is the electrostatic potential at the surface and V_0 the surface potential of the sphere.

The rotational energy is determined within the rigid-body ansatz: $E_{\text{Rot}} = \frac{\hbar^2 l(l+1)}{2I_\perp}$. The surface energy results

*Electronic address: zhanghongfei@impcas.ac.cn

from the effects of the surface tension forces in a half space. When there are nucleons in regard in a neck or a gap between separated fragments an additional term called proximity energy must be added to take into account the effects of the nuclear forces between the close surfaces. This term is essential to describe smoothly the one-body to two-body transition and to obtain reasonable fusion barrier heights. It moves the barrier top to an external position and strongly decreases the pure Coulomb barrier.

$$E_{\text{prox}}(r) = 2\gamma \int_{h_{\text{min}}}^{h_{\text{max}}} \Phi [D(r, h)/b] 2\pi h dh, \quad (8)$$

where h is the distance varying from the neck radius or zero to the height of the neck border. D is the distance between the surfaces in regard and $b = 0.99$ fm the surface width. Φ is the proximity function of Feldmeier [16]. The surface parameter γ is the geometric mean between the surface parameters of the two nuclei or fragments. The combination of the GLDM and of a quasi-molecular shape sequence has allowed to reproduce the fusion barrier heights and radii, the fission and the α and cluster radioactivity data.

For the α emission this very accurate formula simulates the proximity energy [17]:

$$E_{\text{prox}}(r) = (4\pi\gamma)e^{-1.38(r-R_\alpha-R_d)} [0.6584A^{2/3} - (\frac{0.172}{A^{1/3}} + 0.4692A^{1/3})r - 0.02548A^{1/3}r^2 + 0.01762r^3]. \quad (9)$$

To obtain the α decay barrier from the contact point between the nascent α particle and daughter nucleus it is sufficient to add this proximity energy to the Coulomb repulsion.

The half-life of a parent nucleus decaying via α emission is calculated using the WKB barrier penetration probability. In a unified fission model, the decay constant of the α emitter is simply defined as $\lambda = \nu_0 P$. The assault frequency ν_0 has been taken as $\nu_0 = 10^{20} s^{-1}$. The barrier penetrability P is calculated within the action integral

$$P = \exp[-\frac{2}{\hbar} \int_{R_{\text{in}}}^{R_{\text{out}}} \sqrt{2B(r)(E(r) - E(\text{sphere}))}]. \quad (10)$$

The deformation energy (relative to sphere) is small until the rupture point between the fragments [12] and the two following approximations may be used: $R_{\text{in}} = R_d + R_\alpha$ and $B(r) = \mu$ where μ is the reduced mass. R_{out} is simply $e^2 Z_d Z_\alpha / Q_\alpha$. The partial half-life is related to the decay constant λ by $T_{1/2} = \frac{\ln 2}{\lambda}$. The α decay half-lives of the recently produced superheavy nuclei calculated with the GLDM using the experimental Q_α value and without considering the rotational contribution are presented in Table 1. The results agree reasonably with the experimental data indicating that a GLDM taking account the proximity effects, the mass asymmetry, and

an accurate nuclear radius is sufficient to reproduce the α decay potential barriers when the experimental Q_α value is known. The results obtained with the DDM3Y interaction agree with the experimental data as the GLDM predictions and largely better than the VSS calculations. This shows that a double folding potential obtained using M3Y [18] effective interaction supplemented by a zero-range potential for the single-nucleon exchange is very appropriate because its microscopic nature includes many nuclear features, in particular a potential energy surface is inherently embedded in this description. This double agreement shows that the experimental data themselves seem to be consistent. For most nuclei the predictions of the VSS model largely overestimate the half lives. The blocking effect is probably treated too roughly.

The half live of $^{294}118$ is slightly underestimated in the three theoretical calculations possibly due to the neutron submagic number $N = 176$. In Ref. [19], it is also pointed out that for oblate deformed chain of $Z = 112$, the shell closure appears at $N = 176$.

Most of the theoretical half lives using GLDM are slightly smaller than the experimental data. A reason is perhaps that the rotation of the nuclei is neglected in the present calculations. The term $\hbar^2 l(l+1)/(2I_\perp)$ in Eq.(??) represents an additional centrifugal contribution to the barrier which reduces the tunnelling probability and increases the half lives. A second reason is that the shell effects and pairing correlation are not explicitly included in the alpha decay barrier, in spite of their global inclusion in the decay energy Q .

The experimental α decay half-lives are between the close theoretical values given by the GLDM and the ones derived from the VSS formulae. Thus predictions of the α decay half lives with the GLDM and VSS formulae are possible. In the next calculations the experiential Q_α values are taken from the FRDM [20] which reproduces all known experimental data of ground state properties of a large number of nuclei and gives good predictions for nuclei far from the β stability line and the superheavy nucleus region. In Ref. [13] $T_{1/2}$ obtained using the Q_α values given by the Thomas-Fermi model can be found.

α decay half-lives for $Z = 106$ to $Z = 120$ isotopes are shown in Fig.1, the open dots indicating the results of GLDM and the black triangles the ones derived from the VSS formulae.

The half-life values vary from years to microseconds. A narrow window exists between the two predictions for each isotope and the unknown α decay half-lives of SHN may lie in this window assuming that the FRDM Q_α is correct.

The FRDM Q_α value explicitly displays a minimum at the submagic number $N = 162$. It is progressively eroded by the neutron deficiency and the subshell disappears completely from $Z = 115$. This induces a small first hump in the predicted $\log_{10}[T_{1/2}(s)]$ curves till $Z = 114$. Before this first submagic number the half-lives increases rapidly with A from Db to $Z = 114$ isotopes. For the results of

TABLE I: Comparison between experimental α decay half-lives [4] and results obtained with the GLDM, the DDM3Y effective interaction [14] and the VSS formulae.

Parent Z	Nuclei A	Expt. $Q(\text{MeV})$	Expt. $T_{1/2}$	DDM3Y $T_{1/2}$	GLDM $T_{1/2}$	VSS $T_{1/2}$
118	294	11.81 ± 0.06	$1.8^{+7.5}_{-1.3}$ ms	$0.66^{+0.23}_{-0.18}$ ms	$0.15^{+0.05}_{-0.04}$ ms	$0.64^{+0.24}_{-0.18}$ ms
116	293	10.67 ± 0.06	53^{+62}_{-19} ms	206^{+90}_{-61} ms	$22.81^{+10.22}_{-7.06}$ ms	1258^{+557}_{-384} ms
116	292	10.80 ± 0.07	18^{+16}_{-6} ms	39^{+20}_{-13} ms	$10.45^{+5.65}_{-3.45}$ ms	49^{+26}_{-16} ms
116	291	10.89 ± 0.07	$6.3^{+11.6}_{-2.5}$ ms	$60.4^{+30.2}_{-20.1}$ ms	$6.35^{+3.15}_{-2.08}$ ms	$336.4^{+173.1}_{-113.4}$ ms
116	290	11.00 ± 0.08	15^{+26}_{-6} ms	$13.4^{+7.7}_{-5.2}$ ms	$3.47^{+1.99}_{-1.26}$ ms	$15.2^{+9.0}_{-5.6}$ ms
114	289	9.96 ± 0.06	$2.7^{+1.4}_{-0.7}$ s	$3.8^{+1.8}_{-1.2}$ s	$0.52^{+0.25}_{-0.17}$ s	$26.7^{+13.1}_{-8.7}$ s
114	288	10.09 ± 0.07	$0.8^{+0.32}_{-0.18}$ s	$0.67^{+0.37}_{-0.27}$ s	$0.22^{+0.12}_{-0.08}$ s	$0.98^{+0.56}_{-0.40}$ s
114	287	10.16 ± 0.06	$0.51^{+0.18}_{-0.10}$ s	$1.13^{+0.52}_{-0.40}$ s	$0.16^{+0.08}_{-0.05}$ s	$7.24^{+3.43}_{-2.61}$ s
114	286	10.35 ± 0.06	$0.16^{+0.07}_{-0.03}$ s	$0.14^{+0.06}_{-0.04}$ s	$0.05^{+0.02}_{-0.02}$ s	$0.19^{+0.08}_{-0.06}$ s
112	285	9.29 ± 0.06	34^{+17}_{-9} s	75^{+41}_{-26} s	$13.22^{+7.25}_{-4.64}$ s	592^{+323}_{-207} s
112	283	9.67 ± 0.06	$4.0^{+1.3}_{-0.7}$ s	$5.9^{+2.9}_{-2.0}$ s	$0.95^{+0.48}_{-0.32}$ s	$41.3^{+20.9}_{-13.8}$ s
110	279	9.84 ± 0.06	$0.18^{+0.05}_{-0.03}$ s	$0.40^{+0.18}_{-0.13}$ s	$0.08^{+0.04}_{-0.02}$ s	$2.92^{+1.4}_{-0.94}$ s
108	275	9.44 ± 0.07	$0.15^{+0.27}_{-0.06}$ s	$1.09^{+0.73}_{-0.40}$ s	$0.27^{+0.16}_{-0.10}$ s	$8.98^{+5.49}_{-3.38}$ s
106	271	8.65 ± 0.08	$2.4^{+4.3}_{-1.0}$ min	$1.0^{+0.8}_{-0.5}$ min	$0.33^{+0.28}_{-0.16}$ min	$8.6^{+7.3}_{-3.9}$ min




FIG. 1: Comparison between calculated α -decay half-lives of Sg, Hs, Ds, 112, 114, 116, 118 and 120 isotopes using the GLDM and the VSS formulae.

VSS formulae, it seems that the turning point is delayed a little and appears at neutron number $N=163$, for the blocking effect has been magnified and the half lives are overestimated. $N=184$ is always a closure shell. Subclosure or closure shells exist also around $N=176$ for $Z=105$ to $Z=112$. The most stable nuclei should stand about $N=184$ when the proton number is higher than $Z=115$, but stand around $N=176$ for lower Z values.

As a conclusion the half-lives for α -radioactivity have been analyzed in the quasimolecular shape path within a Generalized Liquid Drop Model including the proximity effects between nucleons and the mass and charge asymmetry. The results are in reasonable agreement with the

published experimental data for the alpha decay half-lives of isotopes of charge 112, 114, 116 and 118 and close to the ones derived from the DDM3Y effective interaction. The experimental α decay half lives stand between the GLDM calculations and VSS formulae results and the α decay half-lives of still non-observed superheavy nuclei have been predicted within the GLDM and VSS approaches and Q_α derived from the FRDM.

This work was supported by the Major State Basic Research Development Program in China Under Contract No. G2000077400, the Natural Science Foundation of China (NSFC) under Grant No. 10505016, 10235020, 10235030, 10575119, Knowledge Innovation

Project of the Chinese Academy of Sciences under Grant No KJCX2-SW-N17 and the financial support from DFG

of Germany.

-
- [1] S. Hofmann, G. Münzenberg, *Rev. Mod. Phys.* **72**, 733 (2000).
 - [2] Yu.Ts. Oganessian *et al.*, *Phys. Rev. Lett.* **83**, 3154 (1999); *Nature* **400**, 242 (1999).
 - [3] Yu.Ts. Oganessian *et al.*, JINR Communication D7-2002-287, (2002); Lawrence Livermore National Laboratory Report UCRL-ID-151619 (2003).
 - [4] Yu.Ts. Oganessian *et al.*, *Phys. Rev. C* **70**, 064609 (2004); *Phys. Rev. C* **71**, 029902(E) (2005).
 - [5] P. Armbruster, *Euro. Phys. J. A* **7**, 23 (2000).
 - [6] W. Loveland, K. E. Gregorich, J.B. Patin, D. Peterson, C. Pouki, P. M. Zielinski, and K. Aleklett, *Phys. Rev. C* **66**, 044617 (2002).
 - [7] K.E. Gregorich *et al.*, *Phys. Rev. C* **72**, 014605 (2005).
 - [8] J. Blocki, J. Randrup, W.J. Swiatecki, and C.F. Tsang, *Ann. Phys.* **105**, 427 (1977).
 - [9] G. Royer and B. Remaud, *Nucl. Phys. A* **444**, 477 (1985).
 - [10] G. Royer and K. Zbiri, *Nucl. Phys. A* **697**, 479 (2002).
 - [11] G. Royer, R. Moustabchir, *Nucl. Phys. A* **683**, 182 (2001).
 - [12] G. Royer, *J. Phys. G* **26**, 1149 (2000).
 - [13] G. Royer, K. Zbiri, C. Bonilla, *Nucl. Phys. A* **730**, 355 (2004).
 - [14] P. Roy Chowdhury, C. Samanta and D. N. Basu, *Phys. Rev. C* **73**, 014612 (2006).
 - [15] A. Sobiczewski, Z. Patyk and S. Cwiok, *Phys. Lett. B* **224**, 1 (1989).
 - [16] H. Feldmeier, 12th Summer School on Nuclear Physics, Mikolajki, Poland, 1979.
 - [17] R. Moustabchir and G. Royer, *Nucl. Phys. A* **683**, 266 (2001).
 - [18] G. Bertsch, J. Borysowicz, H. McManus, and W.G. Love, *Nucl. Phys. A* **284**, 399 (1977).
 - [19] X. H. Zhong, L. Li, P. Z. Ning, arXiv: nucl-th/0410083 (2003).
 - [20] P. Möller, J. R. Nix, K. -L. Kratz, *At. Data Nucl. Data Tables* **66**, 131 (1997).