



**HAL**  
open science

## Search for V+A Current in Top Quark Decay in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV

A. Abulencia, J. Adelman, T. Affolder, T. Akimoto, M. G. Albrow, D. Ambrose, S. Amerio, D. Amidei, A. Anastasov, K. Anikeev, et al.

► **To cite this version:**

A. Abulencia, J. Adelman, T. Affolder, T. Akimoto, M. G. Albrow, et al.. Search for V+A Current in Top Quark Decay in  $p\bar{p}$  Collisions at  $\sqrt{s} = 1.96$  TeV. Physical Review Letters, 2007, 98, pp.072001. 10.1103/PhysRevLett.98.072001 . in2p3-00133633

**HAL Id: in2p3-00133633**

<https://in2p3.hal.science/in2p3-00133633v1>

Submitted on 27 Feb 2007

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Search for $V+A$ current in top quark decay in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV

A. Abulencia,<sup>23</sup> J. Adelman,<sup>13</sup> T. Affolder,<sup>10</sup> T. Akimoto,<sup>55</sup> M.G. Albrow,<sup>16</sup> D. Ambrose,<sup>16</sup> S. Amerio,<sup>43</sup> D. Amidei,<sup>34</sup> A. Anastassov,<sup>52</sup> K. Anikeev,<sup>16</sup> A. Annovi,<sup>18</sup> J. Antos,<sup>1</sup> M. Aoki,<sup>55</sup> G. Apollinari,<sup>16</sup> J.-F. Arguin,<sup>33</sup> T. Arisawa,<sup>57</sup> A. Artikov,<sup>14</sup> W. Ashmanskas,<sup>16</sup> A. Attal,<sup>8</sup> F. Azfar,<sup>42</sup> P. Azzi-Bacchetta,<sup>43</sup> P. Azzurri,<sup>46</sup> N. Bacchetta,<sup>43</sup> W. Badgett,<sup>16</sup> A. Barbaro-Galtieri,<sup>28</sup> V.E. Barnes,<sup>48</sup> B.A. Barnett,<sup>24</sup> S. Baroiant,<sup>7</sup> V. Bartsch,<sup>30</sup> G. Bauer,<sup>32</sup> F. Bedeschi,<sup>46</sup> S. Behari,<sup>24</sup> S. Belforte,<sup>54</sup> G. Bellettini,<sup>46</sup> J. Bellinger,<sup>59</sup> A. Belloni,<sup>32</sup> D. Benjamin,<sup>15</sup> A. Beretvas,<sup>16</sup> J. Beringer,<sup>28</sup> T. Berry,<sup>29</sup> A. Bhatti,<sup>50</sup> M. Binkley,<sup>16</sup> D. Bisello,<sup>43</sup> R.E. Blair,<sup>2</sup> C. Blocker,<sup>6</sup> B. Blumenfeld,<sup>24</sup> A. Bocci,<sup>15</sup> A. Bodek,<sup>49</sup> V. Boisvert,<sup>49</sup> G. Bolla,<sup>48</sup> A. Bolshov,<sup>32</sup> D. Bortoletto,<sup>48</sup> J. Boudreau,<sup>47</sup> A. Boveia,<sup>10</sup> B. Brau,<sup>10</sup> L. Brigliadori,<sup>5</sup> C. Bromberg,<sup>35</sup> E. Brubaker,<sup>13</sup> J. Budagov,<sup>14</sup> H.S. Budd,<sup>49</sup> S. Budd,<sup>23</sup> S. Budroni,<sup>46</sup> K. Burkett,<sup>16</sup> G. Busetto,<sup>43</sup> P. Bussey,<sup>20</sup> K. L. Byrum,<sup>2</sup> S. Cabrera,<sup>15</sup> M. Campanelli,<sup>19</sup> M. Campbell,<sup>34</sup> F. Canelli,<sup>16</sup> A. Canepa,<sup>48</sup> S. Carillo,<sup>17</sup> D. Carlsmith,<sup>59</sup> R. Carosi,<sup>46</sup> M. Casarsa,<sup>54</sup> A. Castro,<sup>5</sup> P. Catastini,<sup>46</sup> D. Cauz,<sup>54</sup> M. Cavalli-Sforza,<sup>3</sup> A. Cerri,<sup>28</sup> L. Cerrito,<sup>42</sup> S.H. Chang,<sup>27</sup> Y.C. Chen,<sup>1</sup> M. Chertok,<sup>7</sup> G. Chiarelli,<sup>46</sup> G. Chlachidze,<sup>14</sup> F. Chlebana,<sup>16</sup> I. Cho,<sup>27</sup> K. Cho,<sup>27</sup> D. Chokheli,<sup>14</sup> J.P. Chou,<sup>21</sup> G. Choudalakis,<sup>32</sup> S.H. Chuang,<sup>59</sup> K. Chung,<sup>12</sup> W.H. Chung,<sup>59</sup> Y.S. Chung,<sup>49</sup> M. Ciljak,<sup>46</sup> C.I. Ciobanu,<sup>23</sup> M.A. Ciocci,<sup>46</sup> A. Clark,<sup>19</sup> D. Clark,<sup>6</sup> M. Coca,<sup>15</sup> G. Compostella,<sup>43</sup> M.E. Convery,<sup>50</sup> J. Conway,<sup>7</sup> B. Cooper,<sup>35</sup> K. Copic,<sup>34</sup> M. Cordelli,<sup>18</sup> G. Cortiana,<sup>43</sup> F. Crescioli,<sup>46</sup> C. Cuenca Almenar,<sup>7</sup> J. Cuevas,<sup>11</sup> R. Culbertson,<sup>16</sup> J.C. Cully,<sup>34</sup> D. Cyr,<sup>59</sup> S. DaRonco,<sup>43</sup> S. D'Auria,<sup>20</sup> T. Davies,<sup>20</sup> M. D'Onofrio,<sup>3</sup> D. Dagenhart,<sup>6</sup> P. de Barbaro,<sup>49</sup> S. De Cecco,<sup>51</sup> A. Deisher,<sup>28</sup> G. De Lentdecker,<sup>49</sup> M. Dell'Orso,<sup>46</sup> F. Delli Paoli,<sup>43</sup> L. Demortier,<sup>50</sup> J. Deng,<sup>15</sup> M. Deninno,<sup>5</sup> D. De Pedis,<sup>51</sup> P.F. Derwent,<sup>16</sup> G.P. Di Giovanni,<sup>44</sup> C. Dionisi,<sup>51</sup> B. Di Ruzza,<sup>54</sup> J.R. Dittmann,<sup>4</sup> P. DiTuro,<sup>52</sup> C. Dörr,<sup>25</sup> S. Donati,<sup>46</sup> M. Donega,<sup>19</sup> P. Dong,<sup>8</sup> J. Donini,<sup>43</sup> T. Dorigo,<sup>43</sup> S. Dube,<sup>52</sup> J. Efron,<sup>39</sup> R. Erbacher,<sup>7</sup> D. Errede,<sup>23</sup> S. Errede,<sup>23</sup> R. Eusebi,<sup>16</sup> H.C. Fang,<sup>28</sup> S. Farrington,<sup>29</sup> I. Fedorko,<sup>46</sup> W.T. Fedorko,<sup>13</sup> R.G. Feild,<sup>60</sup> M. Feindt,<sup>25</sup> J.P. Fernandez,<sup>31</sup> R. Field,<sup>17</sup> G. Flanagan,<sup>48</sup> A. Foland,<sup>21</sup> S. Forrester,<sup>7</sup> G.W. Foster,<sup>16</sup> M. Franklin,<sup>21</sup> J.C. Freeman,<sup>28</sup> I. Furic,<sup>13</sup> M. Gallinaro,<sup>50</sup> J. Galyardt,<sup>12</sup> J.E. Garcia,<sup>46</sup> F. Garberson,<sup>10</sup> A.F. Garfinkel,<sup>48</sup> C. Gay,<sup>60</sup> H. Gerberich,<sup>23</sup> D. Gerdes,<sup>34</sup> S. Giagu,<sup>51</sup> P. Giannetti,<sup>46</sup> A. Gibson,<sup>28</sup> K. Gibson,<sup>47</sup> J.L. Gimmell,<sup>49</sup> C. Ginsburg,<sup>16</sup> N. Giokaris,<sup>14</sup> M. Giordani,<sup>54</sup> P. Giromini,<sup>18</sup> M. Giunta,<sup>46</sup> G. Giurgiu,<sup>12</sup> V. Glagolev,<sup>14</sup> D. Glenzinski,<sup>16</sup> M. Gold,<sup>37</sup> N. Goldschmidt,<sup>17</sup> J. Goldstein,<sup>42</sup> G. Gomez,<sup>11</sup> G. Gomez-Ceballos,<sup>11</sup> M. Goncharov,<sup>53</sup> O. González,<sup>31</sup> I. Gorelov,<sup>37</sup> A.T. Goshaw,<sup>15</sup> K. Goulianos,<sup>50</sup> A. Gresele,<sup>43</sup> M. Griffiths,<sup>29</sup> S. Grinstein,<sup>21</sup> C. Grosso-Pilcher,<sup>13</sup> R.C. Group,<sup>17</sup> U. Grundler,<sup>23</sup> J. Guimaraes da Costa,<sup>21</sup> Z. Gunay-Unalan,<sup>35</sup> C. Haber,<sup>28</sup> K. Hahn,<sup>32</sup> S.R. Hahn,<sup>16</sup> E. Halkiadakis,<sup>52</sup> A. Hamilton,<sup>33</sup> B.-Y. Han,<sup>49</sup> J.Y. Han,<sup>49</sup> R. Handler,<sup>59</sup> F. Happacher,<sup>18</sup> K. Hara,<sup>55</sup> M. Hare,<sup>56</sup> S. Harper,<sup>42</sup> R.F. Harr,<sup>58</sup> R.M. Harris,<sup>16</sup> M. Hartz,<sup>47</sup> K. Hatakeyama,<sup>50</sup> J. Hauser,<sup>8</sup> A. Heijboer,<sup>45</sup> B. Heinemann,<sup>29</sup> J. Heinrich,<sup>45</sup> C. Henderson,<sup>32</sup> M. Herndon,<sup>59</sup> J. Heuser,<sup>25</sup> D. Hidas,<sup>15</sup> C.S. Hill,<sup>10</sup> D. Hirschbuehl,<sup>25</sup> A. Hocker,<sup>16</sup> A. Holloway,<sup>21</sup> S. Hou,<sup>1</sup> M. Houlden,<sup>29</sup> S.-C. Hsu,<sup>9</sup> B.T. Huffman,<sup>42</sup> R.E. Hughes,<sup>39</sup> U. Husemann,<sup>60</sup> J. Huston,<sup>35</sup> J. Incandela,<sup>10</sup> G. Introzzi,<sup>46</sup> M. Iori,<sup>51</sup> Y. Ishizawa,<sup>55</sup> A. Ivanov,<sup>7</sup> B. Iyutin,<sup>32</sup> E. James,<sup>16</sup> D. Jang,<sup>52</sup> B. Jayatilaka,<sup>34</sup> D. Jeans,<sup>51</sup> H. Jensen,<sup>16</sup> E.J. Jeon,<sup>27</sup> S. Jindariani,<sup>17</sup> M. Jones,<sup>48</sup> K.K. Joo,<sup>27</sup> S.Y. Jun,<sup>12</sup> J.E. Jung,<sup>27</sup> T.R. Junk,<sup>23</sup> T. Kamon,<sup>53</sup> P.E. Karchin,<sup>58</sup> Y. Kato,<sup>41</sup> Y. Kemp,<sup>25</sup> R. Kephart,<sup>16</sup> U. Kerzel,<sup>25</sup> V. Khotilovich,<sup>53</sup> B. Kilminster,<sup>39</sup> D.H. Kim,<sup>27</sup> H.S. Kim,<sup>27</sup> J.E. Kim,<sup>27</sup> M.J. Kim,<sup>12</sup> S.B. Kim,<sup>27</sup> S.H. Kim,<sup>55</sup> Y.K. Kim,<sup>13</sup> N. Kimura,<sup>55</sup> L. Kirsch,<sup>6</sup> S. Klimentenko,<sup>17</sup> M. Klute,<sup>32</sup> B. Knuteson,<sup>32</sup> B.R. Ko,<sup>15</sup> K. Kondo,<sup>57</sup> D.J. Kong,<sup>27</sup> J. Konigsberg,<sup>17</sup> A. Korytov,<sup>17</sup> A.V. Kotwal,<sup>15</sup> A. Kovalev,<sup>45</sup> A.C. Kraan,<sup>45</sup> J. Kraus,<sup>23</sup> I. Kravchenko,<sup>32</sup> M. Kreps,<sup>25</sup> J. Kroll,<sup>45</sup> N. Krumnack,<sup>4</sup> M. Kruse,<sup>15</sup> V. Krutelyov,<sup>10</sup> T. Kubo,<sup>55</sup> S. E. Kuhlmann,<sup>2</sup> T. Kuhr,<sup>25</sup> Y. Kusakabe,<sup>57</sup> S. Kwang,<sup>13</sup> A.T. Laasanen,<sup>48</sup> S. Lai,<sup>33</sup> S. Lami,<sup>46</sup> S. Lammel,<sup>16</sup> M. Lancaster,<sup>30</sup> R.L. Lander,<sup>7</sup> K. Lannon,<sup>39</sup> A. Lath,<sup>52</sup> G. Latino,<sup>46</sup> I. Lazzizzera,<sup>43</sup> T. LeCompte,<sup>2</sup> J. Lee,<sup>49</sup> J. Lee,<sup>27</sup> Y.J. Lee,<sup>27</sup> S.W. Lee,<sup>53</sup> R. Lefèvre,<sup>3</sup> N. Leonardo,<sup>32</sup> S. Leone,<sup>46</sup> S. Levy,<sup>13</sup> J.D. Lewis,<sup>16</sup> C. Lin,<sup>60</sup> C.S. Lin,<sup>16</sup> M. Lindgren,<sup>16</sup> E. Lipeles,<sup>9</sup> A. Lister,<sup>7</sup> D.O. Litvintsev,<sup>16</sup> T. Liu,<sup>16</sup> N.S. Lockyer,<sup>45</sup> A. Loginov,<sup>36</sup> M. Loreti,<sup>43</sup> P. Loverre,<sup>51</sup> R.-S. Lu,<sup>1</sup> D. Lucchesi,<sup>43</sup> P. Lujan,<sup>28</sup> P. Lukens,<sup>16</sup> G. Lungu,<sup>17</sup> L. Lyons,<sup>42</sup> J. Lys,<sup>28</sup> R. Lysak,<sup>1</sup> E. Lytken,<sup>48</sup> P. Mack,<sup>25</sup> D. MacQueen,<sup>33</sup> R. Madrak,<sup>16</sup> K. Maeshima,<sup>16</sup> K. Makhoul,<sup>32</sup> T. Maki,<sup>22</sup> P. Maksimovic,<sup>24</sup> S. Malde,<sup>42</sup> G. Manca,<sup>29</sup> F. Margaroli,<sup>5</sup> R. Marginean,<sup>16</sup> C. Marino,<sup>25</sup> C.P. Marino,<sup>23</sup> A. Martin,<sup>60</sup> M. Martin,<sup>20</sup> V. Martin,<sup>20</sup> M. Martínez,<sup>3</sup> T. Maruyama,<sup>55</sup> P. Mastrandrea,<sup>51</sup> T. Masubuchi,<sup>55</sup> H. Matsunaga,<sup>55</sup> M.E. Mattson,<sup>58</sup> R. Mazini,<sup>33</sup> P. Mazzanti,<sup>5</sup> K.S. McFarland,<sup>49</sup> P. McIntyre,<sup>53</sup> R. McNulty,<sup>29</sup> A. Mehta,<sup>29</sup> P. Mehtala,<sup>22</sup> S. Menzemer,<sup>11</sup> A. Menzione,<sup>46</sup> P. Merkel,<sup>48</sup> C. Mesropian,<sup>50</sup> A. Messina,<sup>35</sup> T. Miao,<sup>16</sup> N. Miladinovic,<sup>6</sup> J. Miles,<sup>32</sup> R. Miller,<sup>35</sup> C. Mills,<sup>10</sup> M. Milnik,<sup>25</sup> A. Mitra,<sup>1</sup> G. Mitselmakher,<sup>17</sup> A. Miyamoto,<sup>26</sup> S. Moed,<sup>19</sup> N. Moggi,<sup>5</sup> B. Mohr,<sup>8</sup> R. Moore,<sup>16</sup> M. Morello,<sup>46</sup> P. Movilla Fernandez,<sup>28</sup> J. Mülmenstädt,<sup>28</sup>

A. Mukherjee,<sup>16</sup> Th. Muller,<sup>25</sup> R. Mumford,<sup>24</sup> P. Murat,<sup>16</sup> J. Nachtman,<sup>16</sup> A. Nagano,<sup>55</sup> J. Naganoma,<sup>57</sup>  
 I. Nakano,<sup>40</sup> A. Napier,<sup>56</sup> V. Necula,<sup>17</sup> C. Neu,<sup>45</sup> M.S. Neubauer,<sup>9</sup> J. Nielsen,<sup>28</sup> T. Nigmanov,<sup>47</sup> L. Nodulman,<sup>2</sup>  
 O. Norriella,<sup>3</sup> E. Nurse,<sup>30</sup> S.H. Oh,<sup>15</sup> Y.D. Oh,<sup>27</sup> I. Oksuzian,<sup>17</sup> T. Okusawa,<sup>41</sup> R. Oldeman,<sup>29</sup> R. Orava,<sup>22</sup>  
 K. Osterberg,<sup>22</sup> C. Pagliarone,<sup>46</sup> E. Palencia,<sup>11</sup> V. Papadimitriou,<sup>16</sup> A.A. Paramonov,<sup>13</sup> B. Parks,<sup>39</sup> S. Pashapour,<sup>33</sup>  
 J. Patrick,<sup>16</sup> G. Pauletta,<sup>54</sup> M. Paulini,<sup>12</sup> C. Paus,<sup>32</sup> D.E. Pellett,<sup>7</sup> A. Penzo,<sup>54</sup> T.J. Phillips,<sup>15</sup> G. Piacentino,<sup>46</sup>  
 J. Piedra,<sup>44</sup> L. Pinera,<sup>17</sup> K. Pitts,<sup>23</sup> C. Plager,<sup>8</sup> L. Pondrom,<sup>59</sup> X. Portell,<sup>3</sup> O. Poukhov,<sup>14</sup> N. Pounder,<sup>42</sup>  
 F. Prakhoshyn,<sup>14</sup> A. Pronko,<sup>16</sup> J. Proudfoot,<sup>2</sup> F. Ptohos,<sup>18</sup> G. Punzi,<sup>46</sup> J. Pursley,<sup>24</sup> J. Rademacker,<sup>42</sup> A. Rahaman,<sup>47</sup>  
 N. Ranjan,<sup>48</sup> S. Rappoccio,<sup>21</sup> B. Reisert,<sup>16</sup> V. Rekovic,<sup>37</sup> P. Renton,<sup>42</sup> M. Rescigno,<sup>51</sup> S. Richter,<sup>25</sup> F. Rimondi,<sup>5</sup>  
 L. Ristori,<sup>46</sup> A. Robson,<sup>20</sup> T. Rodrigo,<sup>11</sup> E. Rogers,<sup>23</sup> S. Rolli,<sup>56</sup> R. Roser,<sup>16</sup> M. Rossi,<sup>54</sup> R. Rossin,<sup>17</sup> A. Ruiz,<sup>11</sup>  
 J. Russ,<sup>12</sup> V. Rusu,<sup>13</sup> H. Saarikko,<sup>22</sup> S. Sabik,<sup>33</sup> A. Safonov,<sup>53</sup> W.K. Sakumoto,<sup>49</sup> G. Salamanna,<sup>51</sup> O. Saltó,<sup>3</sup>  
 D. Saltzberg,<sup>8</sup> C. Sánchez,<sup>3</sup> L. Santi,<sup>54</sup> S. Sarkar,<sup>51</sup> L. Sartori,<sup>46</sup> K. Sato,<sup>16</sup> P. Savard,<sup>33</sup> A. Savoy-Navarro,<sup>44</sup>  
 T. Scheidle,<sup>25</sup> P. Schlabach,<sup>16</sup> E.E. Schmidt,<sup>16</sup> M.P. Schmidt,<sup>60</sup> M. Schmitt,<sup>38</sup> T. Schwarz,<sup>7</sup> L. Scodellaro,<sup>11</sup>  
 A.L. Scott,<sup>10</sup> A. Scribano,<sup>46</sup> F. Scuri,<sup>46</sup> A. Sedov,<sup>48</sup> S. Seidel,<sup>37</sup> Y. Seiya,<sup>41</sup> A. Semenov,<sup>14</sup> L. Sexton-Kennedy,<sup>16</sup>  
 A. Sfyrla,<sup>19</sup> M.D. Shapiro,<sup>28</sup> T. Shears,<sup>29</sup> P.F. Shepard,<sup>47</sup> D. Sherman,<sup>21</sup> M. Shimojima,<sup>55</sup> M. Shochet,<sup>13</sup>  
 Y. Shon,<sup>59</sup> I. Shreyber,<sup>36</sup> A. Sidoti,<sup>46</sup> P. Sinervo,<sup>33</sup> A. Sisakyan,<sup>14</sup> J. Sjolin,<sup>42</sup> A.J. Slaughter,<sup>16</sup> J. Slaunwhite,<sup>39</sup>  
 K. Sliwa,<sup>56</sup> J.R. Smith,<sup>7</sup> F.D. Snider,<sup>16</sup> R. Snihur,<sup>33</sup> M. Soderberg,<sup>34</sup> A. Soha,<sup>7</sup> S. Somalwar,<sup>52</sup> V. Sorin,<sup>35</sup>  
 J. Spalding,<sup>16</sup> F. Spinella,<sup>46</sup> T. Spreitzer,<sup>33</sup> P. Squillacioti,<sup>46</sup> M. Stanitzki,<sup>60</sup> A. Staveris-Polykalas,<sup>46</sup> R. St. Denis,<sup>20</sup>  
 B. Stelzer,<sup>8</sup> O. Stelzer-Chilton,<sup>42</sup> D. Stentz,<sup>38</sup> J. Strologas,<sup>37</sup> D. Stuart,<sup>10</sup> J.S. Suh,<sup>27</sup> A. Sukhanov,<sup>17</sup> H. Sun,<sup>56</sup>  
 T. Suzuki,<sup>55</sup> A. Taffard,<sup>23</sup> R. Takashima,<sup>40</sup> Y. Takeuchi,<sup>55</sup> K. Takikawa,<sup>55</sup> M. Tanaka,<sup>2</sup> R. Tanaka,<sup>40</sup> M. Tecchio,<sup>34</sup>  
 P.K. Teng,<sup>1</sup> K. Terashi,<sup>50</sup> J. Thom,<sup>16</sup> A.S. Thompson,<sup>20</sup> E. Thomson,<sup>45</sup> P. Tipton,<sup>60</sup> V. Tiwari,<sup>12</sup> S. Tkaczyk,<sup>16</sup>  
 D. Toback,<sup>53</sup> S. Tokar,<sup>14</sup> K. Tollefson,<sup>35</sup> T. Tomura,<sup>55</sup> D. Tonelli,<sup>46</sup> S. Torre,<sup>18</sup> D. Torretta,<sup>16</sup> S. Tourneur,<sup>44</sup>  
 W. Trischuk,<sup>33</sup> R. Tsuchiya,<sup>57</sup> S. Tsuno,<sup>40</sup> N. Turini,<sup>46</sup> F. Ukegawa,<sup>55</sup> T. Unverhau,<sup>20</sup> S. Uozumi,<sup>55</sup> D. Usynin,<sup>45</sup>  
 S. Vallecorsa,<sup>19</sup> N. van Remortel,<sup>22</sup> A. Varganov,<sup>34</sup> E. Vataga,<sup>37</sup> F. Vázquez,<sup>17</sup> G. Velev,<sup>16</sup> G. Veramendi,<sup>23</sup>  
 V. Veszpremi,<sup>48</sup> R. Vidal,<sup>16</sup> I. Vila,<sup>11</sup> R. Vilar,<sup>11</sup> T. Vine,<sup>30</sup> I. Vollrath,<sup>33</sup> I. Volobouev,<sup>28</sup> G. Volpi,<sup>46</sup>  
 F. Würthwein,<sup>9</sup> P. Wagner,<sup>53</sup> R.G. Wagner,<sup>2</sup> R.L. Wagner,<sup>16</sup> J. Wagner,<sup>25</sup> W. Wagner,<sup>25</sup> R. Wallny,<sup>8</sup> S.M. Wang,<sup>1</sup>  
 A. Warburton,<sup>33</sup> S. Waschke,<sup>20</sup> D. Waters,<sup>30</sup> W.C. Wester III,<sup>16</sup> B. Whitehouse,<sup>56</sup> D. Whiteson,<sup>45</sup> A.B. Wicklund,<sup>2</sup>  
 E. Wicklund,<sup>16</sup> G. Williams,<sup>33</sup> H.H. Williams,<sup>45</sup> P. Wilson,<sup>16</sup> B.L. Winer,<sup>39</sup> P. Wittich,<sup>16</sup> S. Wolbers,<sup>16</sup> C. Wolfe,<sup>13</sup>  
 T. Wright,<sup>34</sup> X. Wu,<sup>19</sup> S.M. Wynne,<sup>29</sup> A. Yagil,<sup>16</sup> K. Yamamoto,<sup>41</sup> J. Yamaoka,<sup>52</sup> T. Yamashita,<sup>40</sup> C. Yang,<sup>60</sup>  
 U.K. Yang,<sup>13</sup> Y.C. Yang,<sup>27</sup> W.M. Yao,<sup>28</sup> G.P. Yeh,<sup>16</sup> J. Yoh,<sup>16</sup> K. Yorita,<sup>13</sup> T. Yoshida,<sup>41</sup> G.B. Yu,<sup>49</sup> I. Yu,<sup>27</sup>  
 S.S. Yu,<sup>16</sup> J.C. Yun,<sup>16</sup> L. Zanello,<sup>51</sup> A. Zanetti,<sup>54</sup> I. Zaw,<sup>21</sup> X. Zhang,<sup>23</sup> J. Zhou,<sup>52</sup> and S. Zucchelli<sup>5</sup>

(CDF Collaboration)

<sup>1</sup>*Institute of Physics, Academia Sinica, Taipei, Taiwan 11529, Republic of China*

<sup>2</sup>*Argonne National Laboratory, Argonne, Illinois 60439*

<sup>3</sup>*Institut de Física d'Altes Energies, Universitat Autònoma de Barcelona, E-08193, Bellaterra (Barcelona), Spain*

<sup>4</sup>*Baylor University, Waco, Texas 76798*

<sup>5</sup>*Istituto Nazionale di Fisica Nucleare, University of Bologna, I-40127 Bologna, Italy*

<sup>6</sup>*Brandeis University, Waltham, Massachusetts 02254*

<sup>7</sup>*University of California, Davis, Davis, California 95616*

<sup>8</sup>*University of California, Los Angeles, Los Angeles, California 90024*

<sup>9</sup>*University of California, San Diego, La Jolla, California 92093*

<sup>10</sup>*University of California, Santa Barbara, Santa Barbara, California 93106*

<sup>11</sup>*Instituto de Física de Cantabria, CSIC-University of Cantabria, 39005 Santander, Spain*

<sup>12</sup>*Carnegie Mellon University, Pittsburgh, PA 15213*

<sup>13</sup>*Enrico Fermi Institute, University of Chicago, Chicago, Illinois 60637*

<sup>14</sup>*Joint Institute for Nuclear Research, RU-141980 Dubna, Russia*

<sup>15</sup>*Duke University, Durham, North Carolina 27708*

<sup>16</sup>*Fermi National Accelerator Laboratory, Batavia, Illinois 60510*

<sup>17</sup>*University of Florida, Gainesville, Florida 32611*

<sup>18</sup>*Laboratori Nazionali di Frascati, Istituto Nazionale di Fisica Nucleare, I-00044 Frascati, Italy*

<sup>19</sup>*University of Geneva, CH-1211 Geneva 4, Switzerland*

<sup>20</sup>*Glasgow University, Glasgow G12 8QQ, United Kingdom*

<sup>21</sup>*Harvard University, Cambridge, Massachusetts 02138*

<sup>22</sup>*Division of High Energy Physics, Department of Physics, University of Helsinki and Helsinki Institute of Physics, FIN-00014, Helsinki, Finland*

<sup>23</sup>*University of Illinois, Urbana, Illinois 61801*

<sup>24</sup>*The Johns Hopkins University, Baltimore, Maryland 21218*

- <sup>25</sup>*Institut für Experimentelle Kernphysik, Universität Karlsruhe, 76128 Karlsruhe, Germany*
- <sup>26</sup>*High Energy Accelerator Research Organization (KEK), Tsukuba, Ibaraki 305, Japan*
- <sup>27</sup>*Center for High Energy Physics: Kyungpook National University, Taegu 702-701, Korea; Seoul National University, Seoul 151-742, Korea; and SungKyunKwan University, Suwon 440-746, Korea*
- <sup>28</sup>*Ernest Orlando Lawrence Berkeley National Laboratory, Berkeley, California 94720*
- <sup>29</sup>*University of Liverpool, Liverpool L69 7ZE, United Kingdom*
- <sup>30</sup>*University College London, London WC1E 6BT, United Kingdom*
- <sup>31</sup>*Centro de Investigaciones Energeticas Medioambientales y Tecnologicas, E-28040 Madrid, Spain*
- <sup>32</sup>*Massachusetts Institute of Technology, Cambridge, Massachusetts 02139*
- <sup>33</sup>*Institute of Particle Physics: McGill University, Montréal, Canada H3A 2T8; and University of Toronto, Toronto, Canada M5S 1A7*
- <sup>34</sup>*University of Michigan, Ann Arbor, Michigan 48109*
- <sup>35</sup>*Michigan State University, East Lansing, Michigan 48824*
- <sup>36</sup>*Institution for Theoretical and Experimental Physics, ITEP, Moscow 117259, Russia*
- <sup>37</sup>*University of New Mexico, Albuquerque, New Mexico 87131*
- <sup>38</sup>*Northwestern University, Evanston, Illinois 60208*
- <sup>39</sup>*The Ohio State University, Columbus, Ohio 43210*
- <sup>40</sup>*Okayama University, Okayama 700-8530, Japan*
- <sup>41</sup>*Osaka City University, Osaka 588, Japan*
- <sup>42</sup>*University of Oxford, Oxford OX1 3RH, United Kingdom*
- <sup>43</sup>*University of Padova, Istituto Nazionale di Fisica Nucleare, Sezione di Padova-Trento, I-35131 Padova, Italy*
- <sup>44</sup>*LPNHE, Universite Pierre et Marie Curie/IN2P3-CNRS, UMR7585, Paris, F-75252 France*
- <sup>45</sup>*University of Pennsylvania, Philadelphia, Pennsylvania 19104*
- <sup>46</sup>*Istituto Nazionale di Fisica Nucleare Pisa, Universities of Pisa, Siena and Scuola Normale Superiore, I-56127 Pisa, Italy*
- <sup>47</sup>*University of Pittsburgh, Pittsburgh, Pennsylvania 15260*
- <sup>48</sup>*Purdue University, West Lafayette, Indiana 47907*
- <sup>49</sup>*University of Rochester, Rochester, New York 14627*
- <sup>50</sup>*The Rockefeller University, New York, New York 10021*
- <sup>51</sup>*Istituto Nazionale di Fisica Nucleare, Sezione di Roma 1, University of Rome "La Sapienza," I-00185 Roma, Italy*
- <sup>52</sup>*Rutgers University, Piscataway, New Jersey 08855*
- <sup>53</sup>*Texas A&M University, College Station, Texas 77843*
- <sup>54</sup>*Istituto Nazionale di Fisica Nucleare, University of Trieste/ Udine, Italy*
- <sup>55</sup>*University of Tsukuba, Tsukuba, Ibaraki 305, Japan*
- <sup>56</sup>*Tufts University, Medford, Massachusetts 02155*
- <sup>57</sup>*Waseda University, Tokyo 169, Japan*
- <sup>58</sup>*Wayne State University, Detroit, Michigan 48201*
- <sup>59</sup>*University of Wisconsin, Madison, Wisconsin 53706*
- <sup>60</sup>*Yale University, New Haven, Connecticut 06520*

We report an upper limit on the fraction of  $V + A$  current,  $f_{V+A}$ , in top quark decays, using approximately  $700 \text{ pb}^{-1}$  of  $p\bar{p}$  collisions at  $\sqrt{s}=1.96 \text{ TeV}$  acquired by the upgraded Collider Detector at Fermilab. For the decay  $t \rightarrow Wb \rightarrow \ell\nu b$  (where  $\ell = e$  or  $\mu$ ), the invariant mass of the charged lepton and the bottom quark jet is sensitive to the polarization of the  $W$  boson. We determine  $f_{V+A} = -0.06 \pm 0.25$  given a top quark mass of  $175 \text{ GeV}/c^2$ . We set an upper limit on  $f_{V+A}$  of 0.29 at the 95% confidence level, which represents an improvement by a factor of two on the previous best direct limit.

PACS numbers: 14.65.Ha, 13.88.+e, 12.15.Ji, 12.38.Qk

The decay of the top quark, the most massive fundamental particle observed by experiment [1, 2], is particularly interesting as a direct probe of the charged current weak interaction at the highest energy scale presently available. In the standard model, the spin- $\frac{1}{2}$  top quark decays via the charged current weak interaction to a spin-1  $W^+$  boson and a spin- $\frac{1}{2}$   $b$  quark [3], with a branching fraction above 99% and width  $\Gamma_t = 1.4 \text{ GeV}$  [4] for a top mass of  $175 \text{ GeV}/c^2$ . The lifetime of the top quark,

$\hbar/\Gamma_t \sim 5 \times 10^{-25} \text{ s}$ , is an order of magnitude shorter than the typical strong interaction time-scale for binding of quarks into hadrons,  $\hbar/\Lambda_{QCD} \sim 3 \times 10^{-24} \text{ s}$ . Therefore, the top quark decays before hadronization, and the spin information is directly transferred to the decay products. In the limit  $m_b \rightarrow 0$ , the pure  $V - A$  theory of the weak interaction predicts that the  $b$  quark has left-handed ( $-1/2$ ) polarization (helicity) and the  $W^+$  boson can only have either longitudinal (zero) or left-handed

(−1) polarization. The right-handed (+1) polarization is forbidden. The fraction  $f^0$  of  $W^+$  bosons with longitudinal polarization is predicted at leading order in perturbation theory to be  $f^0 = m_t^2/(2m_W^2 + m_t^2) = 0.70$  [5]. The non-zero  $b$  quark mass and the higher-order QCD and electroweak radiative corrections modify these predictions below the 1% level [6, 7]. However, the presence of non-standard-model couplings in the  $tWb$  vertex could significantly modify the polarization of the top quark decay products [5, 8–10]. Previous results have either been limited by the small statistics of the top quark samples [11–14] or have only set indirect limits [15].

In this Letter, we search for a  $V+A$  current in top quark decay, while assuming that the  $t\bar{t}$  production mechanism is in agreement with the standard model prediction. We further assume the absence of couplings from magnetic moment interactions in the  $tWb$  interaction, so that  $f^0$  is unchanged from 0.70 [5]. Then, the  $V+A$  fraction  $f_{V+A}$  is related to the fraction  $f^+$  of right-handed  $W^+$  bosons by  $f_{V+A} = f^+/(1 - f^0)$ , and the  $V-A$  fraction  $f_{V-A} \equiv 1 - f_{V+A}$  is related to the fraction  $f^-$  of left-handed  $W^+$  bosons by  $f_{V-A} = f^-/(1 - f^0)$ . The  $W^+$  boson polarization can be inferred from the angular distribution of the charged lepton [16] in the decay  $W^+ \rightarrow \ell^+\nu$ ,

$$\frac{1}{\Gamma} \frac{d\Gamma}{d\cos\theta} = \frac{3}{4}(1 - \cos^2\theta)f^0 + \frac{3}{8}(1 - \cos\theta)^2f^- + \frac{3}{8}(1 + \cos\theta)^2f^+,$$

where the angle  $\theta$  is the polar angle of the charged lepton in the rest frame of the  $W^+$  boson. The  $z$ -axis is defined to be the direction of motion of the  $W^+$  boson in the rest frame of the top quark. We use the observable  $M_{\ell b}^2$ , the square of the invariant mass of the charged lepton and the jet from the  $b$  quark, which is related to  $\cos\theta$  by

$$M_{\ell b}^2 \simeq \frac{1}{2}(m_t^2 - m_W^2)(1 + \cos\theta).$$

The relation is exact in the limit  $m_b \rightarrow 0$ .

This search is based on a data set with an integrated luminosity of approximately  $700 \text{ pb}^{-1}$  acquired by the Collider Detector at Fermilab (CDF II)[17] from  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96 \text{ TeV}$ . A 96-layer outer drift chamber [18] reconstructs the trajectories of charged particles and measures their momenta in the region  $|\eta| < 1$  [19]. An eight-layer silicon microstrip detector [20] provides precision tracking information in the region  $|\eta| < 2$  to identify displaced vertices associated with  $b$  hadron decays. The entire tracking volume is located inside a 1.4 T magnetic field. Electromagnetic and hadronic calorimeters measure the energies of particle showers. Drift chambers and scintillation counters provide muon identification outside the calorimeters. Gas Cherenkov counters [21] determine the luminosity. The data are collected with an inclusive lepton trigger that requires an electron (muon) with  $|\eta| < 1$  and  $E_T > 18 \text{ GeV}$  ( $P_T > 18 \text{ GeV}/c$ ) [19].

We study three independent data samples enriched in  $t\bar{t}$  events. Two of the data samples are in the lepton+jets channel, with  $t\bar{t} \rightarrow W^+bW^-\bar{b}$  events where one of the  $W$  bosons decays hadronically and the other leptonically. The lepton+jets event selection requires one isolated lepton with  $E_T > 20 \text{ GeV}$ ,  $\cancel{E}_T > 20 \text{ GeV}$  [19], at least three jets with  $E_T > 15 \text{ GeV}$ , and one or two  $b$ -tagged jets. More details on the selection, the  $b$ -tagging procedure, and the sample composition can be found in Ref. [22]. We model the hard  $t\bar{t}$  process with the Monte Carlo (MC) event generator ALPGEN [23] with CTEQ5L [24] parton densities and PYTHIA [25] for hadronization, under the assumption that the top quark mass is  $175 \text{ GeV}/c^2$ . We simulate the detector response using GEANT [26, 27]. For  $t\bar{t}$  production with  $V-A$  top quark decay, we estimate a selection efficiency, including the branching fraction, of  $\mathcal{A}_{V-A} = 3.4\%(1.2\%)$  for events with one (two)  $b$ -tagged jets. Due to the lower average  $p_T$  of the charged lepton for  $V-A$ , this is a factor 0.92 below the efficiency for  $V+A$ .

For the lepton+jets sample with a single  $b$ -tagged jet, the  $b$ -tagged jet is from the same top quark decay as the charged lepton in approximately half of the  $t\bar{t}$  events. The background  $M_{\ell b}^2$  distribution is a combination of 85%  $W$ +jets, modeled by ALPGEN  $Wb\bar{b}$ , and 15% multi-jet events, modeled by non-isolated lepton+jets data events. Background-dominated data samples with only one jet or only two jets are consistent, in terms of both the rate and the shape of the  $M_{\ell b}^2$  distribution, with our model of the background. In  $695 \text{ pb}^{-1}$ , we observe 304 candidates with a total expected background of  $88 \pm 11$  events.

For the lepton+jets sample with two  $b$ -tagged jets, the two possible  $M_{\ell b}^2$  values of the charged lepton with either the highest or the second highest  $E_T$   $b$ -tagged jet are used to construct a 2-D distribution. In this way, we keep both the correct and incorrect combinations, and account for their correlation. The background is modeled by ALPGEN  $Wb\bar{b}$ ; here the multi-jet background is negligible. Non-uniform binning was applied in the 2-D  $M_{\ell b}^2$  distributions in order to ensure sufficient MC events in each bin. In  $695 \text{ pb}^{-1}$ , we find 75 candidates with a total expected background of  $9 \pm 2$  events.

The third sample is in the dilepton channel, with  $t\bar{t} \rightarrow W^+bW^-\bar{b}$  events where both  $W$  bosons decay leptonically. The dilepton event selection requires two identified leptons with opposite electric charge and  $E_T > 20 \text{ GeV}$ ,  $\cancel{E}_T > 25 \text{ GeV}$ , and at least two jets with  $E_T > 15 \text{ GeV}$ . More details on the selection and the sample composition can be found in Ref. [28]. For  $t\bar{t}$  production with  $V-A$  top quark decay, modeled by ALPGEN as described above, we estimate a selection efficiency, including the branching fraction, of  $\mathcal{A}_{V-A} = 0.72\%$ , a factor 0.88 below the efficiency for  $V+A$ . The two possible  $M_{\ell b}^2$  values for a charged lepton with either the highest or the second highest  $E_T$  jet, assumed to be produced by the fragmentation of the  $b$  quarks, are used to construct a 2-D distribution.

As we can reconstruct  $M_{\ell b}^2$  from the top quark decay and from the anti-top quark decay, we make one entry for each charged lepton. The effect of the correlation between the spins of the top quark and the anti-top quark is negligible here. Again, non-uniform binning in the 2-D  $M_{\ell b}^2$  distributions is applied. The background  $M_{\ell b}^2$  distribution is the combination of three background types: approximately 50% from  $Z/\gamma^* \rightarrow \ell^+\ell^-$  with associated jets, 30% from  $W \rightarrow \ell\nu$  with associated jets where a jet is misidentified as a lepton, and 20% from massive diboson pairs,  $WW/WZ$ . The  $Z/\gamma^*$  and diboson background  $M_{\ell b}^2$  distributions are modeled by ALPGEN. The misidentified lepton background is based on inclusive lepton trigger data where the second lepton is instead a jet (charged particle track) weighted by a probability for misidentification as an electron (muon). A background-dominated data sample with only one jet is consistent, in terms of both the rate and the shape of the  $M_{\ell b}^2$  distribution, with our model of the background. In  $750 \text{ pb}^{-1}$ , we observe 64 candidates (12  $ee$ , 24  $\mu\mu$ , and 28  $e\mu$ ) with a total estimated background of  $20 \pm 4$  events.

The fraction  $f_{V+A}$  is estimated by comparing the  $M_{\ell b}^2$  distribution in data with parent  $M_{\ell b}^2$  distributions for  $t\bar{t}$  production with  $V-A$  top quark decay ( $f_{V+A} = 0.0$ ),  $t\bar{t}$  production with  $V+A$  top quark decay ( $f_{V+A} = 1.0$ ), and backgrounds. A binned log likelihood fit procedure is used to extract the parameter of interest,  $f_{V+A}$ . We represent the imperfectly known accepted background cross section for each sample,  $\sigma_{bg}$ , and the  $t\bar{t}$  cross section [29, 30],  $\sigma_{t\bar{t}}$ , by nuisance parameters. The analytic expression for the likelihood for each sample,

$$\mathcal{L} = \left[ \prod_{i=0}^N P(n_i, \mu_i) \right] \times G(\sigma_{bg}, \delta\sigma_{bg}) \times G(\sigma_{t\bar{t}}, \delta\sigma_{t\bar{t}}), \quad (1)$$

is the product over all  $N$  bins in  $M_{\ell b}^2$  of the Poisson probabilities of observing  $n_i$  entries in a given bin  $i$ , where the average expected bin content is  $\mu_i$ , and the Gaussian constraints on the estimated background and the predicted  $t\bar{t}$  production cross sections, as shown in Table I. The  $\mu_i$  are given by:

$$\mu_i = N^{data} \left[ x_{V+A} \hat{T}_{V+A}^i + x_{V-A} \hat{T}_{V-A}^i + x_{bg} \hat{T}_{bg}^i \right], \quad (2)$$

$$x_{V+A} = \frac{f_{V+A} \mathcal{A}_{V+A} \sigma_{t\bar{t}}}{\sigma_{bg} + \sigma_{t\bar{t}} [\mathcal{A}_{V+A} f_{V+A} + \mathcal{A}_{V-A} (1 - f_{V+A})]}, \quad (3)$$

$$x_{V-A} = \frac{(1 - f_{V+A}) \mathcal{A}_{V-A} \sigma_{t\bar{t}}}{\sigma_{bg} + \sigma_{t\bar{t}} [\mathcal{A}_{V+A} f_{V+A} + \mathcal{A}_{V-A} (1 - f_{V+A})]}, \quad (4)$$

$$x_{bg} = \frac{\sigma_{bg}}{\sigma_{bg} + \sigma_{t\bar{t}} [\mathcal{A}_{V+A} f_{V+A} + \mathcal{A}_{V-A} (1 - f_{V+A})]}. \quad (5)$$

Here,  $N^{data}$  is the total number of observed events for the sample. The  $x_{V+A}$ ,  $x_{V-A}$ , and  $x_{bg}$  are the fractions of  $t\bar{t}$  production with  $V+A$  top quark decay,  $t\bar{t}$  production with  $V-A$  top quark decay, and background, respectively. The

TABLE I: The input values for the nuisance parameters, and the values from the best fit to the combined samples.

Nuisance parameter	Input (pb)	Fit (pb)
$\sigma_{t\bar{t}}$	$6.7 \pm 1.0$	$7.3 \pm 0.9$
$\sigma_{bg}$ lepton+jets 1 $b$ -tag	$0.156 \pm 0.017$	$0.154 \pm 0.016$
$\sigma_{bg}$ lepton+jets 2 $b$ -tag	$0.013 \pm 0.002$	$0.013 \pm 0.002$
$\sigma_{bg}$ dilepton	$0.026 \pm 0.006$	$0.022 \pm 0.006$

$\hat{T}_{V+A}^i$ ,  $\hat{T}_{V-A}^i$ , and  $\hat{T}_{bg}^i$  are the probabilities for an event to occupy bin  $i$  of the corresponding  $M_{\ell b}^2$  distribution. Note that  $\sum_i \hat{T}^i = 1.0$ . The combined likelihood is the product of the likelihoods of the three samples, but with one common Gaussian constraint on the  $t\bar{t}$  cross section.

The robustness of the fitting procedure has been tested with MC simulated experiments. For a given experiment and a particular sample, the number of observed data events is distributed in three categories ( $t\bar{t}$  production with  $V+A$  top quark decay,  $t\bar{t}$  production with  $V-A$  top quark decay, and background) according to their expected fractions from Equations 3-5. For each category, the events are generated from the relevant  $M_{\ell b}^2$  parent distribution. The hypotheses that  $f_{V+A} = 0.0, 0.1, \dots, 1.0$  are studied for 2000 experiments for all samples combined, as well for the three samples separately. In all cases, the fit is unbiased and stable. An expected statistical uncertainty of 0.22 on  $f_{V+A}$  is found for the combined case, while for the separate samples we found 0.35 for lepton+jets events with one  $b$ -tagged jet, 0.41 for lepton+jets events with two  $b$ -tagged jets, 0.27 for all lepton+jets events, and 0.46 for dilepton events. In all cases, this uncertainty includes a small component ( $\leq 0.02$ ) due to the uncertainties on the background and  $t\bar{t}$  cross sections.

The maximum likelihood fit to the lepton+jets sample yields a value of  $f_{V+A} = 0.21 \pm 0.28$ , with  $f_{V+A} = 0.16 \pm 0.36$  for the subset of events with one  $b$ -tagged jet and  $f_{V+A} = 0.28 \pm 0.44$  for the subset of events with two  $b$ -tagged jets. For the dilepton sample, we obtain  $f_{V+A} = -0.64 \pm 0.37$ . The probability to obtain a value smaller than the dilepton result is 10% for the hypothesis  $f_{V+A} = 0$ . The lepton+jets and dilepton results are compatible at about 1.8 standard deviations.

The estimates of the systematic uncertainties on the measured value for  $f_{V+A}$  are shown in Table II for all samples combined. The leading sources of systematic uncertainty arise from uncertainties on the measured jet energy [31], the background shape and normalization, and limited MC statistics. We determine all systematic uncertainties by performing MC simulated experiments in which the systematic parameter in question is varied and the resulting simulated data are fit to the default parent distributions. All systematic uncertainties are evaluated

TABLE II: The systematic uncertainties on the measurement of  $f_{V+A}$  for all samples combined.

Source	Uncertainty
Jet energy	0.10
Background modeling	0.04
MC statistics	0.04
Initial/Final state QCD radiation	0.02
Multiple $p\bar{p}$ interactions	0.02
$b$ -tag efficiency ( $E_T$ )	0.02
MC generator	0.01
Parton densities	0.01
Total	0.12




FIG. 1: Likelihood distribution (see Equation 1) for the lepton+jets and dilepton data samples separately and combined.

at  $f_{V+A} = 0$ ; we find the dominant uncertainty is insensitive to the value of  $f_{V+A}$ .

Many of the above sources of systematic uncertainty also contribute significantly to the systematic uncertainty on the measurement of the top quark mass [32]. We choose to quote our result at a top quark mass value of  $175 \text{ GeV}/c^2$ , where the dependence of the measurement (upper limit) for  $f_{V+A}$  is  $\pm 0.07 (\pm 0.06)$  for a  $\mp 2 \text{ GeV}/c^2$  shift in top quark mass.

Combining all samples, the result for the fraction of  $V+A$  current in top quark decay is

$$f_{V+A} = -0.06 \pm 0.22 \text{ (stat.)} \pm 0.12 \text{ (syst.)}.$$

This value is in agreement with the standard model. Table I summarizes the fitted values for the nuisance parameters. The likelihood distribution is shown in Fig. 1 for the combined sample as well as for each individual sample. The good agreement in the  $M_{\ell b}^2$  distribution between data and the best fit result for  $f_{V+A}$  from the combined sample is shown in Figs. 2, 3, and 4, where the highest bins also contain overflow entries. For comparison,  $f_{V+A} = 1.00$  is also shown.




FIG. 2: The  $M_{\ell b}^2$  distribution for the lepton+jets sample with a single  $b$ -tagged jet. For  $f_{V+A} = -0.06$ , the  $\chi^2$  probability is 69%.

In the absence of a signal, we evaluate an upper limit on  $f_{V+A}$  using a Bayesian approach. The profile likelihood function is first determined as a function of  $f_{V+A}$ , multiplied by a prior flat between 0.0 and 1.0, and normalized to yield the posterior distribution for  $f_{V+A}$ . The upper limit at 95% confidence level (CL) is formed by integrating the posterior from zero to the value of  $f_{V+A}$  that yields 0.95 for the integral. We verified that this approach yields proper frequentist coverage for  $f_{V+A} \leq 0.3$ ; a small correction, derived by the Neyman construction, would be applied to any upper limit greater than 0.3 to restore coverage in the region  $f_{V+A} > 0.3$ . From MC simulated experiments for the standard model case where  $f_{V+A} = 0$ , the median upper limit is 0.38 and 68% of experiments set an upper limit between 0.2 and 0.6.

Combining all samples, we set an upper limit on the fraction of  $V+A$  current in top quark decay of

$$f_{V+A} < 0.29 \text{ at 95\% CL.}$$

This is an improvement by a factor of two on the previous best direct limit [11]. Converting the results for  $f_{V+A}$  to the fraction of right-handed  $W^+$  bosons, we obtain  $f^+ = -0.02 \pm 0.07 \text{ (stat.)} \pm 0.04 \text{ (syst.)}$  and  $f^+ < 0.09$  at 95% CL.

We thank the Fermilab staff and the technical staffs of the participating institutions for their vital contributions. This work was supported by the U.S. Department of Energy and National Science Foundation; the Italian Istituto Nazionale di Fisica Nucleare; the Ministry of Education, Culture, Sports, Science and Technology of Japan; the Natural Sciences and Engineering Research Council of Canada; the National Science Council of the Republic of China; the Swiss National Science Foundation; the A.P. Sloan Foundation; the Bundesministerium für Bildung und Forschung, Germany; the Korean Science and Engineering Foundation and the Korean Research Foundation; the Particle Physics and Astronomy Research Council and the Royal Society, UK; the Institut




FIG. 3: The invariant mass squared of the charged lepton and the highest  $E_T$   $b$ -tagged jet,  $M^2_{lb1}$ , in five regions of the invariant mass squared of the charged lepton and the second highest  $E_T$   $b$ -tagged jet,  $M^2_{lb2}$ , for the lepton+jets sample with two  $b$ -tagged jets. For  $f_{V+A} = -0.06$ , the  $\chi^2$  probabilities are 92%, 3%, 10%, 18%, and 47% in order of increasing  $M^2_{lb2}$ .




FIG. 4: The invariant mass squared of a charged lepton and the highest  $E_T$  jet,  $M^2_{lb1}$ , in four regions of the invariant mass squared of the charged lepton and the second highest  $E_T$  jet,  $M^2_{lb2}$ , for the dilepton sample. There are two entries per event, one for each lepton. For  $f_{V+A} = -0.06$ , the  $\chi^2$  probabilities are 58%, 14%, 54%, and 51% in order of increasing  $M^2_{lb2}$ .

National de Physique Nucleaire et Physique des Particules/CNRS; the Russian Foundation for Basic Research; the Comisi3n Interministerial de Ciencia y Tecnologia, Spain; the European Community's Human Potential Programme under contract HPRN-CT-2002-00292; and the Academy of Finland.

[1] F. Abe et al. (CDF Collaboration), Phys. Rev. Lett. **74**, 2626 (1995).  
 [2] S. Abachi et al. (DØ Collaboration), Phys. Rev. Lett. **74**, 2632 (1995).

[3] Charge conjugate decays, with replacement of left-handed by right-handed polarization, are implicit.  
 [4] M. Jezabek and J. H. Kuhn, Nucl. Phys. B **314**, 1 (1989).  
 [5] G. L. Kane, G. A. Ladinsky, and C. P. Yuan, Phys. Rev. D **45**, 124 (1992).  
 [6] M. Fischer, S. Groote, J. G. Korner, and M. C. Mauser, Phys. Rev. D **63**, 031501(R) (2001).  
 [7] H. S. Do, S. Groote, J. G. Korner, and M. C. Mauser, Phys. Rev. D **67**, 091501(R) (2003).  
 [8] M. A. B. Beg, R. V. Budny, R. Mohapatra, and A. Sirlin, Phys. Rev. Lett. **38**, 1252 (1977).  
 [9] G. Triantaphyllou, J. Phys. G **26**, 99 (2000).  
 [10] D. Choudhury, T. M. P. Tait, and C. E. M. Wagner, Phys. Rev. D **65**, 053002 (2002).  
 [11] D. Acosta et al. (CDF Collaboration), Phys. Rev. D **71**, 031101(R) (2005).  
 [12] A. Abulencia et al. (CDF Collaboration), Phys. Rev. D **73**, 111103(R) (2006).  
 [13] V.M. Abazov et al. (DØ Collaboration), Phys. Lett. B **617**, 1 (2005).  
 [14] V.M. Abazov et al. (DØ Collaboration), Phys. Rev. D **72**, 011104(R) (2005).  
 [15] F. Larios, M. A. Perez, and C. P. Yuan, Phys. Lett. B **457**, 334 (1999).  
 [16] We use lepton to denote an electron or a muon.  
 [17] D. Acosta et al. (CDF Collaboration), Phys. Rev. D **71**, 032001 (2005).  
 [18] A. Affolder et al., Nucl. Instrum. Methods A **526**, 249 (2004).  
 [19] We use a cylindrical coordinate system where the  $z$  axis is along the proton beam direction and  $\theta$  is the polar angle. Pseudorapidity is  $\eta = -\ln \tan(\theta/2)$ , while transverse momentum is  $p_T = |p| \sin \theta$ , and transverse energy is  $E_T = E \sin \theta$ . Missing transverse energy,  $\cancel{E}_T$ , is defined as the magnitude of  $-\sum_i E_T^i \hat{n}_i$ , where  $\hat{n}_i$  is the unit vector in the azimuthal plane that points from the beam line to the  $i^{th}$  calorimeter tower.  
 [20] A. Sill, Nucl. Instrum. Methods A **447**, 1 (2000).  
 [21] D. Acosta et al., Nucl. Instrum. Methods A **494**, 57 (2002).  
 [22] D. Acosta et al. (CDF Collaboration), Phys. Rev. D **71**, 052003 (2005).  
 [23] M. L. Mangano et al., J. High Energy Phys. **0307**, 001 (2003).  
 [24] H. L. Lai et al. (CTEQ Collaboration), Eur. Phys. J. C **12**, 375 (2000).  
 [25] T. Sjostrand et al., Comput. Phys. Commun. **135**, 238 (2001).  
 [26] E. Gerchtein and M. Paulini, eConf **C0303241**, TUMT005 (2003).  
 [27] R. Brun et al., CERN-DD/EE/84-1 (1987).  
 [28] D. Acosta et al. (CDF Collaboration), Phys. Rev. Lett. **93**, 142001 (2004).  
 [29] M. Cacciari et al., J. High Energy Phys. **404**, 68 (2004).  
 [30] N. Kidonakis and R. Vogt, Phys. Rev. D **68**, 114014 (2003).  
 [31] A. Bhatti et al., Nucl. Instrum. Methods A **566**, 375 (2006).  
 [32] W. M. Yao et al., J. Phys. G **33**, 1 (2006).