



Quarkonium polarization in Pb-Pb and pp collisions with ALICE

Y. Ding

► To cite this version:

Y. Ding. Quarkonium polarization in Pb-Pb and pp collisions with ALICE. 41st International Conference on High Energy Physics (ICHEP 2022), Jul 2022, Bologna, Italy. in2p3-04916047


HAL Id: in2p3-04916047

<https://in2p3.hal.science/in2p3-04916047v1>

Submitted on 28 Jan 2025

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.


L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



ICHEP 2022
BOLOGNA



Université Claude Bernard Lyon 1



Quarkonium polarization in Pb—Pb and pp collisions with ALICE


Yanchun Ding^{1,2} for the ALICE Collaboration

Central China Normal University¹


Institut de Physique des Deux Infinis de Lyon²

6th - 13th July, 2022





Introduction




ALICE



Polarization:

- Defined as the particle spin-alignment with respect to a chosen direction
- Measured via anisotropies in the decay products angular distributions

$$W(\cos \theta, \varphi) \propto \frac{1}{3 + \lambda_\theta} (1 + \lambda_\theta \cos^2 \theta + \lambda_\varphi \sin^2 \theta \sin 2\varphi + \lambda_{\theta\varphi} \sin 2\theta \cos \varphi)$$



GOAL: obtain λ_θ , λ_φ and $\lambda_{\theta\varphi}$

$$(\lambda_\theta, \lambda_\varphi, \lambda_{\theta\varphi}) = (0, 0, 0)$$

No polarization

$$(\lambda_\theta, \lambda_\varphi, \lambda_{\theta\varphi}) = (+1, 0, 0)$$


Pure transverse polarization

$$(\lambda_\theta, \lambda_\varphi, \lambda_{\theta\varphi}) = (-1, 0, 0)$$

Pure longitudinal polarization




How to define polarization axis




Reference frames:

- 👉 **Helicity (HE)**: the direction of quarkonium in the center-of-mass frame
- 👉 **Collins-Soper (CS)**: the bisector of the angle between the direction of one beam and the opposite of the other beam in the quarkonium rest frame





Physics motivation



ALICE

Polarization in pp collisions: constrain quarkonium production mechanisms

Theoretically



J/ ψ :

- ↳ NLO NRQCD ==> transverse polarization in Helicity frame
- ↳ NLO CSM ==> longitudinal polarization in Helicity frame



M. Butenschoen *et al.*, *Phys. Rev. Lett.* 108 (2012) 172002



Υ :

- ↳ NLO NRQCD ==> no significant polarization for $\Upsilon(1S)$ and $\Upsilon(2S)$ states, but a strong transverse polarization for $\Upsilon(3S)$ at high p_T
- ↳ ICEM (k_T -factorization approach) ==> transverse polarization and no significant differences in polarization among the $\Upsilon(nS)$ states




B. Gong *et al.*, *Phys. Rev. Lett.* 112 (2014) 3, 032001



C. Vincent *et al.*, *Phys. Rev. D* 99 (2019) 3, 034007



No sizeable polarization is observed for the existing quarkonium polarization measurements in pp collisions



Polarization in Pb—Pb collisions

🔔 Potential difference w.r.t pp collisions

👉 In central collisions

- ↳ Modification of prompt J/ψ feed-down fractions due to $\psi(2S)$ and χ_c suppression in the QGP

$$J/\psi^{\text{Prompt}} : (60\%)^{\text{Direct}} + (30\%)^{\chi_c} + (10\%)^{\psi(2S)}$$

📎 J.-P. Lansberg, *Phys. Rept.* 889 (2020)

- ↳ Contribution from charmonium (re)generation

👉 In non-central collisions

- ↳ Large angular momentum due to the medium rotation is predicted


📎 F. Becattini et al., *Phys. Rev. C* 77, 024906

- ↳ Huge magnetic field is expected


📎 D. E. Kharzeev et al., *Nucl. Phys. A* 803 (2008) 227-253



A Large Ion Collider Experiment



ALICE



A dedicated heavy-ion experiment at the LHC



Muon Spectrometer ($-4 < \eta < -2.5$)

- I. Absorbers: efficiently dump π , K and low-momentum muons
- II. Tracking system + III. Dipole magnet: muon track reconstruction, muon momentum and its electric charge measurement
- IV. Trigger system: muon PID and unlike sign dimuon trigger (for the quarkonium analyses)



Inclusive quarkonium detection down to $p_T = 0$

The other quarkonium talks from ALICE:

Raphaelle Marie Bailhache 7th July, 2022

Theraa M A Tork 8th July, 2022


Maurice Louis Coquet 9th July, 2022

Himanshu Sharma 9th July, 2022

Biswarup Paul 9th July, 2022



A Large Ion Collider Experiment



ALICE




A dedicated heavy-ion experiment at the LHC


☞ V. Silicon Pixel Detector

- ↳ Vertex reconstruction
- ↳ Event Plane determination


☞ VI. V0 Detectors

- ↳ Event trigger
- ↳ Centrality determination
- ↳ Background rejection
- ↳ Event plane determination






$\Upsilon(1S)$ polarization in pp collisions at $\sqrt{s} = 13$ TeV




ALICE





🔔 First ALICE $\Upsilon(1S)$ polarization measurement in pp collisions

👉 λ_θ , λ_φ and $\lambda_{\theta\varphi}$ consistent with zero within uncertainties in both HE and CS frames



$\Upsilon(1S)$ polarization in pp collisions



🔔 First **ALICE** $\Upsilon(1S)$ polarization measurement in pp collisions

👉 λ_θ , λ_φ and $\lambda_{\theta\varphi}$ consistent with zero within uncertainties in both **HE** and **CS** frames

👉 Compatible with **Pb—Pb** results

📎 *Phys. Lett. B* 815 (2021) 136146

👉 Good agreement with **LHCb** pp data at $\sqrt{s} = 8$ TeV, in a similar rapidity range, within the large experimental uncertainties


📎 **LHCb** Collaboration, *JHEP* 12 (2017) 110

👉 Qualitatively described by **NLO NRQCD** calculations

📎 M. Butenschoen et al., *Phys. Rev. Lett.* 108 (2012) 172002



J/ ψ polarization in pp and Pb—Pb collisions



ALICE

ALICE measured J/ ψ polarization in Pb—Pb collisions

- All polarization parameters are close to zero within uncertainties
- ↳ λ_θ shows a maximum 2σ deviation w.r.t zero in both HE and CS frames for $2 < p_T < 4$ GeV/c

- Compatible with ALICE results in pp collisions within uncertainties

EPJC 78 (2018) 562

- 3σ difference w.r.t LHCb in pp collisions in HE frame


LHCb Collaboration, EPJC 73 (2013) 11

→ Difference due to suppression/regeneration effects in Pb—Pb w.r.t pp collisions?

→ What is the role of the angular momentum (\vec{L}) and the magnetic fields (\vec{B})?




Phys. Lett. B 815 (2021) 136146





J/ ψ polarization as a function of the event plane



ALICE

J/ ψ polarization in Pb—Pb collisions measured using another reference frame

Reference frame:


- ☞ Event-plane based frame (EP): axis orthogonal to the EP in the collision center-of-mass frame


- ☞ EP normal to \vec{B} and \vec{L}

☞ Significant spin alignment observed for light vector mesons (K^{*0} , Φ) Phys. Rev. Lett 125 (2020) 012301


Heavy-quark pair production:

- ☞ Occurs early in the collision ($t \sim 0.1 \text{ fm}/c$)
- ☞ Experiences both the short living \vec{B} and the \vec{L} of the rotating medium






J/ ψ polarization as a function of EP in Pb—Pb collisions



ALICE

arXiv:2204.10171




First measurement of J/ψ polarization w.r.t the EP




Centrality dependence



Significant polarization (3.5σ) observed in 40-
60% for $2 < p_T < 6 \text{ GeV}/c$




J/ ψ polarization as a function of EP in Pb–Pb collisions



arXiv:2204.10171

ALICE



🔔 First measurement of J/ ψ polarization w.r.t the EP

👉 p_T dependence

↳ Significant deviation (3.9σ) at low p_T ($2 < p_T < 4$ GeV/c) for 30–50%

↳ Similar to light-flavor hadrons (K^{*0} , Φ): maximum polarization at low p_T for semicentral collisions, but:

↳ Smaller absolute polarization

$$|\lambda_\theta^{J/\psi}| < |\lambda_\theta^\Phi| < |\lambda_\theta^{K^{*0}}|$$

↳ Different sign of the deviation


$$\lambda_\theta^{J/\psi} > 0, \lambda_\theta^{\Phi, K^{*0}} < 0$$

📎 Phys. Rev. Lett 125 (2020) 012301

- 📌 Different production mechanisms for J/ ψ and light-flavor hadrons in nuclear collisions
- 📌 Different rapidity range for the two measurements



Summary



ALICE

🔔 First ALICE $\Upsilon(1S)$ polarization measurement in pp collisions

- 👉 All polarization parameters are compatible with zero in both **HE** and **CS** frames

🔔 First J/ψ polarization measurement as a function of the event plane in Pb—Pb collisions


- 👉 Significant deviation w.r.t zero is observed for λ_θ in semicentral collisions at low p_T

Outlook




➡ Theoretical description of vector meson polarization in heavy-ion collisions is needed

➡ More precise measurements can be expected from large data sample with LHC Run 3




Happy ending




ALICE






$\Upsilon(1S)$ polarization in pp collisions: analysis strategy



ALICE


Signal extraction

- Raw number of $\Upsilon(1S)$ obtained by fitting the dimuon invariant mass distribution



Polarization parameters determination

- λ_θ , λ_φ and $\lambda_{\theta\varphi}$ extracted by fitting to the $A \times \varepsilon$ -corrected $\Upsilon(1S)$ angular distributions in both frames simultaneously



Acceptance x efficiency correction

- Corrected number of $\Upsilon(1S)$ evaluation based on a MC simulation



J/ ψ polarization vs EP in Pb—Pb collisions: analysis strategy



ALICE

arXiv:2204.10171


Signal extraction


- ↳ **Reweighting** applied at the dimuon candidate level with a 2-dimensions (p_T , $\cos \theta$) $A \times \varepsilon$ map
- ↳ Fitting the corrected dimuon invariant mass distribution for the extraction of the raw yield

Polarization parameters determination


- ↳ Fitting the corrected angular distributions and extracting the polarization parameters

- ➡ **Cross check:** λ_θ compatible with zero when evaluated w.r.t a **random EP**






$\Upsilon(1S)$ polarization in Pb—Pb collisions



ALICE



🔔 First **ALICE** $\Upsilon(1S)$ polarization measurement in pp collisions


👉 λ_θ , λ_ϕ and $\lambda_{\theta\phi}$ consistent with zero within uncertainties in both **HE** and **CS** frames

👉 Compatible with **Pb—Pb** results

📎 Phys. Lett. B 815 (2021) 136146




Light flavor hadrons (K^{*0} , Φ) polarization



ALICE

EP



PP


RP

- 🔔 ρ_{00} measurement for light flavor hadrons in Pb—Pb collisions at $\sqrt{s_{\text{NN}}} = 2.76 \text{ TeV}$ and in pp collisions at $\sqrt{s} = 13 \text{ TeV}$
- 👉 p_{T} dependence
 - ↳ $\rho_{00} < 1/3$ for K^{*0} and Φ at low p_{T} (smaller central value for K^{*0}) in Pb—Pb collisions
 - ↳ $\rho_{00} \sim 1/3$ for:
 - ↳ $p_{\text{T}}^{K^{*0}} > 2 \text{ GeV}/c$ and $p_{\text{T}}^{\Phi} > 0.8 \text{ GeV}/c$
 - ↳ A random event plane (RP)
 - ↳ K^{*0} and Φ in pp collisions

➡ Zero spin hadron K_S^0 : no spin alignment is observed



Light flavor hadrons (K^{*0} , Φ) polarization



ALICE


🔔 ρ_{00} measurement for light flavor hadrons in Pb—Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV and in pp

collisions at $\sqrt{s} = 13$ TeV

👉 Centrality dependence

↳ ρ_{00} deviates w.r.t 1/3 at low p_T in **semi-central collisions**


↳ No centrality dependence of ρ_{00} at high p_T



ALI-PUB-520392




Charmed mesons polarization




ALICE

- 🔔 Charmed vector meson (D^{*+}) polarization crucial to complete the picture in HICs
- 🔔 D^{*+} polarization in pp collisions at $\sqrt{s} = 13$ TeV
 - 👉 ρ_{00} spin matrix element (1/3 means no polarization)
 - 👉 Prompt D^{*+} ($c \rightarrow D^{*+}$) unpolarized
 - 👉 Non-zero polarization for non-prompt D^{*+} ($b \rightarrow D^{*+}$)
 - 👉 Both well predicted by PYTHIA 8 + EVTGEN





Angular distribution



ALICE

$$W(\cos \theta) \propto \frac{1}{3 + \lambda_\theta} (1 + \lambda_\theta \cos^2 \theta)$$


$$W(\varphi) \propto 1 + \frac{2\lambda_\varphi}{3 + \lambda_\theta} \cos 2\varphi$$

$$W(\tilde{\varphi}) \propto 1 + \frac{\sqrt{2}\lambda_{\theta\varphi}}{3 + \lambda_\theta} \cos 2\tilde{\varphi}$$

$\tilde{\varphi} = \varphi - 3\pi/4, \cos \theta < 0$
 $\tilde{\varphi} = \varphi - \pi/4, \cos \theta > 0$



J/ ψ polarization in theory



ALICE

Phys. Rev. Lett. 108 (2012) 172002

