



HAL
open science

Physics Prospects for ALICE in Run 5 and Beyond

Antonio Uras

► **To cite this version:**

Antonio Uras. Physics Prospects for ALICE in Run 5 and Beyond. 9th Large Hadron Collider Physics Conference (LHCP2021), Jun 2021, Paris, France. in2p3-04916405

HAL Id: in2p3-04916405

<https://in2p3.hal.science/in2p3-04916405v1>

Submitted on 28 Jan 2025

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

D. Longhi et al. LHC2021



LHCP2021

The Ninth Annual Conference on Large Hadron Collider Physics



7-12 June 2021 ~~Paris (France), Sorbonne Université (IN2P3/CNRS, IRFU/CEA)~~

Physics Prospects for ALICE in Run 5 and Beyond



Antonio Uras
IP2I - Lyon – CNRS/IN2P3

on behalf of the ALICE Collaboration




ALICE 3: new dedicated heavy-ion experiment at the LHC, replacing ALICE starting of Run 5: hadronization mechanisms in the medium, QGP transport properties, access to the pre-equilibrium phase

- **Discussed at the heavy-ion town meeting** (CERN, Oct 2018)

<https://arxiv.org/abs/1902.01211>

- **Expression of Interest** submitted as input to the European Particle Physics Strategy Update (Granada, May 2019)

In this presentation >>> Selected material from the preliminary discussions on the ALICE 3 physics goals and scientific opportunities, and the detector concept




see also talks by C. Lippman, G. Contin, S. Bufalino

Precision temperature measurement with uncertainties comparable to low-energy experiments?


Effects of chiral symmetry restoration, predicted by QCD, can be studied at the LHC at vanishing μ_B

- Effect on ρ - a_1 mixing on the dilepton mass spectrum above ϕ peak
- In-medium broadening of narrow vector resonances?

Measurement of pre-equilibrium dileptons through multi-differential (p_T , flow, polarization, DCA) measurements: **fireball chronometer**



❖ **Electric conductivity, or electric charge diffusion coefficient:** response of an equilibrated relativistic gas of electrically charged particles, upon the influence of a small, static, electric field



❖ **Lower and upper limits of thermal dilepton production spectra connected to QGP conductivity:** spectra can be exploited to constrain predictions on the QGP electric conductivity

Precise data are needed to challenge theoretical models on the estimation of the diffusion coefficients of the strongly interacting QGP

Phys. Rev. D 93, 096012 (2016)

❖ **Soft photons** ($p_T^\gamma \ll p_T^{\text{hadrons}} \approx 300\text{-}500 \text{ MeV}$) can be produced at any stage of hadronic collisions, with no specific constraints in their number by conservation laws

❖ **Low's theorem:** QCD prediction providing a precise relation between very soft photon and inclusive hadron production

$$\frac{dN_\gamma}{d^3k} = \frac{\alpha}{2\pi k_0} \int d^3p_1 d^3p_2 d^3p_3 \dots d^3p_N \sum_{i,j=1}^N \eta_i \eta_j e_i e_j \frac{-(p_i \cdot p_j)}{(p_i \cdot k)(p_j \cdot k)} \frac{dN_{\text{hadrons}}}{d^3p_1 d^3p_2 d^3p_3 \dots d^3p_N}$$

❖ **Soft photon puzzle:** nearly every measurement shows factor 2–5 enhancement w.r.t. Low's theorem predictions. **Proposed explanations:** cold quark-gluon plasma, quark synchrotron radiation, string fragmentation. Handle to investigate fundamental non-perturbative properties of QCD




Ultra-light converter-tracker + calorimeter at forward η should allow measuring soft photons down to $p_T \approx 10 \text{ MeV}$ (possibly exploiting HBT analysis techniques)

- ❖ In heavy-ion collisions, large increase of multi-HF baryons (≈ 1000) expected via coalescence with charm quarks from different hard scatterings ($N_c \approx 100$ in central Pb-Pb)

Multi-charm baryons are almost pure coalescence particles: (potentially) large discrimination power on the role of the various hadronization mechanisms

Ω_{cc} and Ω_{ccc} not yet observed. Ω_{ccc} may only be accessible in heavy-ion collisions

Establish clean **reconstruction of decay cascade**, exploiting state-of-the-art vertexing and tracking





$$\Omega_{ccc}^{++} \rightarrow \Omega_{cc}^+ + \pi^+$$

$$\Omega_{cc}^+ \rightarrow \Omega_c^0 + \pi^+$$


$$\Omega_c^0 \rightarrow \Omega^- + \pi^+$$

$$\Omega^- \rightarrow \Lambda + K^-$$

$$\Lambda \rightarrow p + \pi^-$$



Quarkonium Measurements



Complete spectroscopy of states in the QGP → study **direct** exclusive quarkonium production by subtraction of the feed-down components

χ_c states:


- Binding energy in between J/ψ and $\psi(2S)$
- Sizable feed-down contribution to J/ψ
- **Most promising decay mode: $\chi_c \rightarrow J/\psi \gamma$ (γ measured with calorimetry and/or pair conversion)**

Pseudoscalar η_c states


- Similar behaviour in the QGP w.r.t. vector states e.g. J/ψ
- Factorisation approach + heavy-quark spin symmetry assumption allows for the simultaneous treatment of J/ψ and $\eta_c(1S)$

Hadrons with more than 3 valence quarks for which we don't have a complete understanding of their nature: e.g. X(3872)


- **Detailed and differential study in heavy-ion collisions proposed as a tool to indirectly constrain its nature:** production yield in the dense QCD environment could be largely influenced by its inner structure



- If the mystery of its nature is addressed by the end of Run 4 we will have a **new, tuned tool to study HF hadronization in the QGP**



Low- p_T reach crucial for a full characterization of the hadronization mechanism




- ❖ **Direct measurement of the fragmentation patterns of charmed/beauty mesons and baryons**
- ❖ **Jets provide energy and direction scale for the fragmentation process: proxy for initial HF quark direction and energy**


- ❖ **Studying the fragmentation shower of quarkonium and open HF inside jets in AA collisions: new insights into the properties of in-medium propagation of quarkonium states inside the QGP**



Low- p_T reach needed for a complete picture of the fragmentation functions




Double Parton Scattering: Quarkonia and Open HF



Measurements of the production of quarkonia “in association” with another final state particle

Double parton scattering: two independent scatterings in one pp/pA collision

- ❖ **Powerful probe to study** factorization of hard processes in hadronic collisions, and transverse parton densities in nucleons and nuclei
- ❖ **DPS events characterized by large pseudorapidity gap between the two hadrons:** → At large $\Delta\eta$ pure DPS “environment”



Phys. Rev. D 90, 111101(R)

Dark Photons: hypothetical extra-U(1) gauge bosons, motivated by:


- Antiproton spectrum and positron excess in cosmic ray observations
- Muon anomalous magnetic moment

Possible channels in ALICE 3:

- Meson decays such as π^0 , η , ϕ Dalitz decays, D^{*0} decays, radiative J/ψ and Υ decays
- **Final-state radiation, Drell-Yan, thermal rad. for $M > 1$ GeV**
- Displaced searches ($M < 20$ MeV)

Requirements for ALICE 3


- Good electron ID capability for wide momentum range (low momenta from π^0 Dalitz decays to high momenta from DY and thermal dielectrons)
- High-rate capability and in-bunch pileup separation + good vertexing to separate thermal dielectrons and HF pairs






Ultra-peripheral heavy-ion collisions (UPC): clean environment + huge $Z^4 \approx 5 \cdot 10^7$
 enhanced gamma+gamma rate w.r.t. pp

❖ **Searches of BSM particle coupling predominantly to photons:** modifications of the light-by-light scattering rates from virtual corrections from heavy particles (magnetic monopoles, vector-like fermions, dark sector particles)

❖ **Precision measurements of EM couplings of SM particles:** anomalous magnetic moment ($g-2$) of the tau



Challenge for ALICE 3: acceptance for tau and light-by-light scattering down to low p_T ?





State-of-the art experiment optimized for “soft” physics from pp to Pb-Pb:

- Tracking and vertexing accuracy down to zero p_T
- Complete suite of PID detectors
- Large rapidity coverage
- Extreme acquisition rates for soft, untriggerable probes
- Unique vertexing capabilities

Important deadlines:



- Open ALICE 3 workshop in September/October
- End 2021: submission of the Lol to the LHCC



ALICE 3 will be equipped with a $\approx 100 \text{ m}^2$ tracker based on large, bended MAPS sensors

Retractable layers (IRIS) under study:

Getting closer to the interaction point during stable beam ($R = 0.5, 1.2, 2.5 \text{ cm}$)



Ultra-light tracker:

$\approx 0.05 \% X_0$ vertexing layers

$\approx 0.5 \% X_0$ tracking layers




Large acceptance: $|\eta| < 4.0$, full azimuth down to very low p_T

Great potential for charm measurements

ALICE 3 will be equipped with a $\approx 100 \text{ m}^2$ tracker based on large, bended MAPS sensors

Retractable layers (IRIS) under study:

Getting closer to the interaction point during stable beam ($R = 0.5, 1.2, 2.5 \text{ cm}$)



Ultra-light tracker:

$\approx 0.05 \% X_0$ vertexing layers


$\approx 0.5 \% X_0$ tracking layers

Large acceptance: $|\eta| < 4.0$, full azimuth down to very low p_T


Great potential for charm measurements

Several PID options and technologies under investigation (performance, costs...)

TOF detector outside the tracker:
 $R \approx 100\text{cm} (+20\text{cm?}), \sigma = 20\text{ ps}$



Aerogel-RICH detector:
 $R \approx 120\text{ cm}, 50\text{ ps time res.}$



- ❖ **Barrel ECAL:** $E(\text{photons}) \approx 0.1\text{ GeV} - 10/20\text{ GeV}$, $R \approx 130\text{ cm}$, PbWO_4 crystals
- ❖ **Forward conversion tracker:** photons down to $p_T \approx 10\text{ MeV}$, $2.5 < \eta < 5$
 → **Unique possibility to test soft theorems for photon production at the LHC**
- ❖ **MuonID:** muons down to $p_T \approx 1.5\text{ GeV}$ at $\eta = 0$, $R \approx 160\text{ cm}$, GEM technology
 → **Unique possibility for charmonia down to zero p_T at $\eta = 0$ at the LHC**
- ❖ **Pixel Shower Detector** (radiator + high-granularity pixels) to improve electron ID

- ❖ **ALICE is preparing a next-generation heavy-ion experiment for LHC Run 5 and beyond:** access to novel measurements of electromagnetic and hadronic probes of the QGP at very low momenta

- ❖ **Physics goal: measurements** inaccessible in LHC Run 3+4 because of limitations in detector performance or available luminosity:
 - High-precision measurements of dielectron production
 - New measurements of quarkonium states, multi-charm baryons and exotica
 - Searches for signals of new physics beyond Standard Model

- ❖ **Performance studies and detector R&D plans are ongoing**
 - Physics cases and detector options explored by dedicated working groups, discussed in internal ALICE workshops
 - First "open" workshop in September/October 2021, LoI to be submitted end of 2021

Backup Slides