



HAL
open science

Détecter la matière noire sub-GeV

J. Gascon, A. Letessier Selvon, D. Santos

► **To cite this version:**

J. Gascon, A. Letessier Selvon, D. Santos. Détecter la matière noire sub-GeV. Séminaire Thématique "Physique des neutrinos et matière noire", Oct 2019, TALENCE, France. in2p3-04926814

HAL Id: in2p3-04926814

<https://in2p3.hal.science/in2p3-04926814v1>

Submitted on 3 Feb 2025

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Détecter la matière noire sub-GeV

***Gas Detectors for Direct and Directional Dark Matter Detection
and Axion-Like Particle exploration***

***The direct search for Dark Matter in the keV/c² to GeV/c²
range through its interaction with nucleons or electrons using
solid-state cryogenic detectors***

Direct detection of dark matter in the eV-to-GeV mass range

J. Gascon (IP2I),
pour A. Letessier-Selvon (LPNHE) et D. Santos (LPSC)
+ autres co-auteurs de ces trois contributions

■ **Physics of direct DM searches in the eV-GeV range**

- Physics arguments of the 3 contributions based on discussions already documented elsewhere
- Present summary largely based on: *Dark Sectors 2016 Workshop: Community Report* (arXiv:1608.08632)

See also updates in « US Cosmic Visions: New Ideas in Dark Matter 2017: Community Report » arXiv:1707.04591

3 contributions :

- Gas & directional detectors
- Skipper-CCD
- Cryogenic detectors

**More details on
detector technologies
in GT08 contributions**

Physics motivation

Dark Sectors 2016 Workshop: Community Report (arXiv:1608.08632):



- *"In recent years it has been shown that many production mechanisms that explain the observed DM relic abundance, point to light, sub-GeV DM, below the reach of standard direct detection techniques. It is thus crucial to develop direct-detection strategies for such low DM masses."*
- $m_\chi > \text{eV}$ if fermion (Galactic DM halo = Fermi gas)
- $m_\chi > 3 \text{ keV}$ if thermal equilibrium with SM particles (Ly- α forest)
- $E_{\text{NR}} \leq 2(\mu_{\chi,N} v_\chi)^2 / m_N$, with $v_\chi \sim 10^{-3} c$
- For a Si nucleus and $m_\chi \sim 100 \text{ MeV}$, $E_{\text{NR}} \sim 1 \text{ eV}$
- Scattering on electron favored as m_χ approaches m_e
- Complementarity with searches at beam dumps & colliders

DM models in eV-GeV range



- **Freeze-out DM:** relic density obtained from thermal equilibrium of SM+SM \rightarrow DM+DM, but via **Dark Sector mediators** instead of SM mediators. *e.g. Dark Photon, with small couplings to standard γ*
- **Asymmetric DM:** final relic abundance of DM from particle/antiparticle asymmetry in the DM sector. **Natural mass \sim GeV**, but keV-GeV possible depending on DM sector
- **Freeze-in DM:** *no thermal equilibrium of DM with SM:* instead, SM particles slowly annihilate or decay into DM, via DS
- **SIMPs** (strongly interacting) thermally coupled to SM, relic abundance set by (3 \rightarrow 2 freeze-out): **natural sub-GeV mass + high DD rates**
- **Elastically Decoupling Relic** : DM relic abundance determined by its inelastic scattering with SM: MeV to $<$ GeV and DD signal possible
- **Bosonic DM (can be $<$ 1 keV):** QCD axion, axion-like particles, and the dark photon itself. **DM entirely absorbed in detector in DD experiments. Axions produced in the sun** can also be detected.

Detection signals

- Fermion: scattering off nuclei or electrons




- Bosons: Primakoff effect (ALPs), photoabsorption (dark photons)



Scattering: materials vs mass & energy scale

US Cosmic Visions: New Ideas in Dark Matter 2017: Community Report [arXiv:1707.04591]





See also: K. Zurek, CERN-th colloquium 03.08.16

Electron vs nucleon scattering



Dark Sectors 2016 Workshop: Community Report (arXiv:1608.08632):

- \sim eV to few eV thresholds possible in semiconductor/scintillators: sensitivity to MeV DM via electron scattering
- meV thresholds in (future) superconducting detectors \rightarrow keV DM
- Nucleon scattering adversely affected by *kinematics* and by *quenching*


arXiv:1608.08632



Inelastic scattering on atoms: Migdal effect





- Similar to bremsstrahlung effect
- Inelastic scattering on atom (nucleus+e⁻'s): transferred energy not limited by two-body kinematics
- Increases sensitivity to scattering of **Nuclear Recoils** (energy boost), but relatively rare process ($<10^{-3}$)
- **Electron signal** (no quenching)
- *Calculation only: an experimental confirmation would be welcomed*



Electron searches

- Two cases: Dark Photon heavier (left) or lighter (right) than DM particle
- Most interesting models (on or above green/orange/blue lines) can be probed using a 1 kg-y exposure of a detector with zero background + single-electron sensitivity
- If dark sector DM subdominant: need larger bkg-free exposure



Goals and Challenges

Dark Sectors 2016 Workshop: Community Report (arXiv:1608.08632):

- Low threshold
 - DM kinetic energy $\sim (1 \text{ eV}) * (M_{\text{DM}}/1 \text{ MeV})$
- Backgrounds
 - Bkg usually associated to $>1 \text{ GeV}$ searches (γ / neutrons from internal+external radioactivity, cosmic rays, activation) are less important than detector-specific bkg (e.g. dark current)
- Scaling up exposure
 - Low threshold usually associated with small detectors
 - Internal/external radioactivity becomes an issue again
- Signal discrimination and background model
- Material

Gas Detectors for Direct and Directional Dark Matter
Detection and Axion-Like Particle exploration:
NEWS-G and gas TPC

GAS DETECTORS


Gas detectors: NEWS-G

- NEWS-G (New Experiment for Wimps detection with a Sphere filled with Gas)
- Direct search for very-low mass WIMPs, from 0.1 to 10 GeV.
 - 1.4 diameter, single-channel readout
 - Sub-keV energy threshold
 - Fiducialisation via pulse shape
 - Background rejection by pulse shape analysis
 - Up to 10 bars: ten's of kg with various light gas targets: H, He, Ne.
- First tests at LSM: summer/autumn 2019.



NEWS-G@LSM physics reach

- Installation of NEWS-G at SNOLAB is scheduled November 2019
- Physics reach: sub-GeV with light gas targets




- + Quenching measurements @ LPSC

Gas detector: TPC for axion

- MIMAC: gaseous micro-TPC with pixelized Micromega readout, designed to test the directionality of an eventual DM signal
- Proposal for an application for ALP DM in the mass window from 100 eV up to 20 keV
- *Kaluza-Klein Axion-Like particles: long lifetimes, detection of its decays into two identical photons (no strong electric field needed).*
- New generation of micro-TPCs with 100 eV energy threshold and a high spatial 3D resolution in a large active volume.
- Background reduced by the many discrimination observables offered by this technique.
- Matrix of many individual bi-chamber modules (with low individual background rates) for a total of 2 m³.


Physics reach of gas TPC for axions

- Rate of KK-axion decays detected MIMAC active volume (300 mbar, $^{40}\text{Ar} + 5\% \text{C}_4\text{H}_{10}$) as a function of ALP mass. Strongly increase when the flux from those trapped by the sun (red) is added to the direct flux from the Sun (blue).



Sensitivity to trapped axions

- MIMAC 1-y exclusion region (blue) vs axion density on Earth compared with the XMASS exclusion (red) and NEWS-G exclusion (green). MIMAC should reach the Kaluza-Klein-axion model (cf dot: not yet excluded).



Direct detection of dark matter in the eV-to-GeV mass
Range

SKIPPER-CCDS

Skipper CCDs

- Improved signal/noise from repeated measurement of the charge (wrt pedestal)




Figure 3 : a) CCD output signal for one pixel illustrating the principle of CDS (R: reset, S: charge transfer). b) Skipper vs. conventional readout: during the same time of a conventional readout, the pixel charge is measured N times in skipper mode. The effect of a low-frequency noise waveform is much smaller in skipper mode because the waveform changes very little during the short CDS sequence.

- 0.07 electron resolution already achieved





Figure 4: Pixel charge distribution of a DAMIC-M skipper CCD. Peaks corresponding to pixels with zero, one and two charges are clearly distinguished. A Gaussian fit to the charge distribution yields a resolution of 0.07 electrons [23].

Skipper-CCD

- Skipper-CCD's already produced world leading limits for sub-GeV DM (SENSEI and DAMIC@SNOLAB)
- Arrays of 100 g (SENSEI) and 1 kg (DAMIC-M @LSM) under construction.
- Future project: 10 kg experiment to probe sub-GeV DM even if it makes only a small fraction of the halo density

DM-electron limits (heavy/light Dark Mediator) + dark photon limits, based on 1 and 2 electron signal, assuming negligible background



Advantages:

- **Very low leakage current**, $< 10^{-21}$ A/cm² at 140 K
- Excellent **spatial resolution**
- Background identification and rejection, **3D reconstruction for surface events**, follow up of **radioactive decay** chains using **spatial correlation of events**.

GT08: Toward a 10-kg skipper-CCD sensor

- Must develop new fabrication method for the large number of skipper-CCDs required by the 10 kg experiment. New foundry needed for future fabrication: opportunity to migrate into the more modern CMOS technology.
- Low noise readout for 10 kg system: several thousand channels, comparable in size with that of the largest CCD cameras ever built for astronomy, but with more stringent noise requirements.
- Needed background more than an order of magnitude lower than planned for SuperCDMS at SNOLAB (goal of zero ionization events in the 2-10 e⁻ signal range)

- Involvement in DAMIC-M @LSM; low-radioactivity.
- LSST+DAMIC-M: expertise in read-out and CMOS technology that are required for development of future 10 kg array
- Simulation and data analysis
- Strong connexion with US partners (Fermilab, Chicago, Washington, PNNL): US project for 10 kg

The direct search for Dark Matter in the keV/c² to GeV/c² range through its interaction with nucleons or electrons using solid-state cryogenic detectors

CRYOGENIC DETECTORS

SubGeV program (2020-2025)

1 kg array of Ge heat-and-ionization detectors with

- Full event-by-event ER/NR discrimination above 50 eV when operated at low bias (<10 V) – *NR searches of 0.1-1 GeV DM*
- Single-electron thresholds when operated at high bias (100 V and more) – *ER searches 0.5-10⁵ MeV DM*
- Statistical separation (and comprehensive study) of backgrounds from the comparison of the data recorded at HV/LV

30 g Ge units with

- $\sigma=10$ eV phonon resolution (18 eV already achieved)
- $\sigma=20$ eV_{ee} ionization resolution (via HEMT preamp)
- 100 V achieved on large 800 g Ge, 400V on 200g



SubGeV program (2020-2025)

30 g Ge units with


- $\sigma = 10$ eV phonon resolution (18 eV already achieved)
- $\sigma = 20$ eV_{ee} ion. resolution (devel. of cold HEMT preamp)

100 V achieved on large 800 g Ge; 400V on 200g

$\sigma = 0.6$ electron already achieved on 30 g @ 70V



• RED30 : 33 g Ge Al electrodes, NTD thermal sensor



Instrumentation challenges

Cf contributions to cryogenic detectors in GT08

- Dual (phonon+ionization) readout important for the study & control of instrumental backgrounds (heat-only, dark current due to trapping)
- R&D to combine high-resolution thermal readout with a tag on one charge (or more) using superconducting thin film sensors inspired by the Superconducting Single Photon Detectors (SSPD)
- Beyond 2025: extend cryogenic detector expertise to superconducting metal absorbers able to reach few meV energy thresholds and study their electromagnetic background suppression capabilities.

Solid-state cryogenics detectors @ in2p3

- Expertise in design, production and operation of solid-state cryogenic detectors: phonon sensor developments, low-noise charge readout, control of leakage currents.
- Very strong synergy with two other major projects using solid-state cryogenic detectors: Ricochet-CENNS and CUPID-Mo/CROSS double beta decay experiments
- Close relationship with the French company CryoConcept - best designer of ultra-low vibration dry dilution refrigerators
- Long standing experience and investment in a very-low background experiments at the LSM.
- Leadership position in EDELWEISS
- Collaboration with US and Canadian partners in SuperCDMS.

CONCLUSIONS

Conclusions

- Strong physics motivation & technological opportunity to explore eV-GeV range of DM mass with direct detection experiments
- Rapid advances in development of new detectors dedicated to that range – *Technological developments central issue*
 - Detector backgrounds, scalability
 - Contributions to GT08
- In France:
 - Gas detectors: NEW—G @SNOLAB, Mimac/ALPS
 - Skipper-CCD detectors with DAMIC-M @ LSM (1 kg) and future 10 kg
 - Cryogenic detectors @ LSM (heat+LN boost, single-electron tag)
- Note: LSM used in all 3 projects
 - Deepest lab in Europe, easy access: excellent platform for these detector developments, and <100 kg experiments.

BACKUPS

Scattering: materials vs mass & energy scale

Dark Sectors 2016 Workshop: Community Report (arXiv:1608.08632)




FIG. 11: Materials that could be used to probe sub-GeV DM, down to keV masses, by *scattering* off electrons [e⁻] or nuclei [N]. Certain DM candidates, which can instead be *absorbed* by bound electrons in these materials, could be probed down to meV masses (not shown). Adapted from [173].

See also: K. Zurek, CERN-th colloquium 03.08.16

Dark Sectors 2016 Workshop: Community Report (arXiv:1608.08632):

Material	$m_{\text{DM,th}}$ (theoretical)	Technology	Challenges	(Optimistic) Timescale
Noble liquids (Xe, Ar)	few MeV	two-phase TPC	dark counts	existing
Semiconductors (Si, Ge)	$\sim 0.1 - 1$ MeV	CCDs & Calorimeter	dark counts (?)	$\sim 1 - 2$ years
Scintillators (GaAs, NaI, CsI)	$\sim 0.5 - 1$ MeV	Calorimeter: $\sigma_E \sim 0.2$ eV	sensitivity & afterglow (?)	$\lesssim 5$ years
Superconductors (Al)	~ 1 keV	Calorimeter: $\sigma_E \sim 1$ meV	sensitivity & unknown backgrounds	$\sim 10 - 15$ years
Superfluid He (NR)	~ 1 MeV	Calorimeter: $\sigma_E \sim 1$ eV	sensitivity & unknown backgrounds	$\lesssim 5$ years
Bond Breaking	\sim few MeV	color centers	sensitivity & unknown backgrounds	$\lesssim 5$ years
Superfluid He (2-excitation)	~ 1 keV	Calorimeter $\sigma_E \sim 10$ meV	sensitivity & unknown backgrounds	$\sim 5 - 10$ years
2D-targets (graphene)	few MeV	based on PTOLEMY	low exposure, unknown backgrounds	$\sim 5 - 10$ years

TABLE III: Material, theoretical mass threshold, required technology to achieve lowest mass threshold, potential or known challenges, and optimistic timescales. All materials and techniques, besides two-phase TPCs in noble liquids, still need to be demonstrated to have sensitivity to sub-GeV DM. Most timescales are thus only illustrative of the time needed to study their feasibility. Materials/techniques sensitive to DM-electron (DM-nucleus) interactions are at the top (bottom) of the table.