



HAL
open science

Impact of intracellular radionuclide distribution in Targeted Alpha Therapy: a Monte-Carlo biophysical study in 3D multicellular model

V Levrague, M E Alcocer-Àvila, L Maigne, M Beuve, E Testa, R Delorme

► **To cite this version:**

V Levrague, M E Alcocer-Àvila, L Maigne, M Beuve, E Testa, et al.. Impact of intracellular radionuclide distribution in Targeted Alpha Therapy: a Monte-Carlo biophysical study in 3D multicellular model. IV Geant4 International User Conference at the physics-medicine-biology frontier, Oct 2022, Naples, Italy. in2p3-04954493

HAL Id: in2p3-04954493

<https://in2p3.hal.science/in2p3-04954493v1>

Submitted on 18 Feb 2025

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**“ Impact of intracellular radionuclide
distribution in Targeted Alpha Therapy : a
Monte-Carlo biophysical study in 3D
multicellular model “**

V. Levrague¹, M.E. Alcocer-Àvila², L. Maigne³, M. Beuve², E. Testa², R. Delorme¹

¹LPSC, Université Grenoble Alpes, 38026 Grenoble Cedex, France

²IP2I, Université Claude Bernard Lyon 1, 69622 Villeurbanne Cedex, France

³Université Clermont Auvergne, CNRS/IN2P3, LPC, 63000 Clermont-Ferrand, France

Background: To understand and predict the therapeutic efficiency of targeted alpha therapy, nano/micro-dosimetry are needed by considering the very heterogeneous dose deposition at cell level. The objective of this study is to evaluate theoretically the importance of radionuclide cell internalization on relevant dosimetric and biological endpoints.

Material and Methods: The treatment with ²¹¹At of mono-cells or microtumors was simulated. The Monte Carlo CPOP [1] code, based on Geant4, was used to generate realistic deformable 3D multicellular geometries, and was adapted to generate the radionuclide source distributions in various internalization cases (membrane, cytoplasm, cytoplasm + nucleus and nucleus only) and collect the physical outputs needed for the biophysical calculations. Physical absorbed doses in cell nuclei of ovarian cancer cell line OVCAR-3, have been calculated, in addition to therapeutic indexes like Tumor Control Probability (TCP), using the biophysical model NanOx [2], which considers in the current version that the cell nucleus is the unique sensitive volume.

The impact of cell packing, tumor size, alphas energy and radionuclide daughter diffusion after fixation were studied, under the hypothesis that all cells are labeled with alpha particles.

Preliminary results: With 42 alpha particles per cell, radionuclide cell internalization has very little impact on physical dose and TCP, except if radionuclides are internalized in the nucleus. However, when the number of alphas per cell is reduced below 10, TCP decreases below 1. For instance, with 5 alphas per cell, TCP were equal to 0.06, 0.16 and 0.66 when the sources were distributed, respectively, in membrane, cytoplasm and nucleus only.

[1] Maigne, L., et al. Phys. Med. 89 (2021) 41-50

[2] Cunha, M. et al. Phys. Med. Biol. 62 (2017) 12-48